

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 219/00	A1	(11) International Publication Number: WO 97/26245
		(43) International Publication Date: 24 July 1997 (24.07.97)

(21) International Application Number: PCT/US97/00015

(22) International Filing Date: 15 January 1997 (15.01.97)

(30) Priority Data:
08/585,090 16 January 1996 (16.01.96) US
08/683,927 19 July 1996 (19.07.96) US

(60) Parent Applications or Grants

(63) Related by Continuation

US 08/585,090 (CIP)
Filed on 16 January 1996 (16.01.96)
US 08/683,927 (CIP)
Filed on 19 July 1996 (19.07.96)

(71) Applicant (for all designated States except US): LUMIGEN, INC. [US/US]; 24485 W. Ten Mile Road, Southfield, MI 48034 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): AKHAVAN-TAFTI, Hashem [US/US]; 2107 Johns Drive, Brighton, MI 48116 (US). ARGHAVANI, Zahra [IR/US]; 2107 Johns Drive, Brighton, MI 48116 (US). DESILVA, Renuka [LK/US]; 40628 Delta Drive, Northville, MI 48167 (US).

(74) Agents: BOLAND, Mark et al.; Sughrue, Mion, Zinn, MacPeak & Seas, Suite 800, 2100 Pennsylvania Avenue, N.W., Washington, DC 20037-3202 (US).

(81) Designated States: AU, CA, CN, JP, KR, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

(54) Title: COMPOUNDS, COMPOSITIONS AND METHODS FOR GENERATING CHEMILUMINESCENCE WITH PHOSPHATASE ENZYMES

(57) Abstract

Novel heterocyclic compounds which generate chemiluminescence on reaction with a phosphatase enzyme are provided as well as a process for their preparation and intermediates useful therein. The compounds comprise a nitrogen, oxygen or sulfur-containing heterocyclic ring system bearing an exocyclic carbon-carbon double bond. The double bond is further substituted at the distal carbon with a phosphate group and an oxygen or sulfur atom-containing group. Novel compositions further comprising a cationic aromatic compound (CAC) in addition to the heterocyclic phosphate compound are provided. The addition of the CAC in the composition greatly increases the production of chemiluminescence and provides improved detection sensitivity. Compositions further comprising an anionic surfactant and a non-ionic surfactant provide additional improvements in detection sensitivity. The novel chemiluminescent compounds and compositions are useful in methods for producing light and in assays for phosphatase enzymes and enzyme inhibitors and in assays employing enzyme-labeled specific binding pairs.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

COMPOUNDS, COMPOSITIONS AND METHODS FOR GENERATING
CHEMILUMINESCENCE WITH PHOSPHATASE ENZYMES

CROSS REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of
applicant's co-pending U.S. Applications Serial Nos.
5 08/585,090 filed on January 16, 1996 and 08/683,927 filed
on July 19, 1996.

FIELD OF THE INVENTION

The present invention relates to chemiluminescent
10 compounds which react with phosphatase enzymes to generate
light. In particular, the present invention relates to
chemiluminescent compounds containing a heterocyclic ring
group and an enol phosphate group which react with oxygen
upon removal of a phosphate group with a phosphatase enzyme
15 to produce an enolate which reacts with to produce
chemiluminescence and a carbonyl compound.

The present invention further relates to compositions
for generating chemiluminescence by reaction with a
phosphatase enzyme. Chemiluminescent compositions comprise
20 a first compound containing a heterocyclic ring group and
an enol phosphate group and a second compound which acts to
increase light production from the reaction of the
phosphate compound with a phosphatase enzyme. The present
invention relates to methods for generating light or
25 chemiluminescence by the reaction of a phosphatase enzyme
with a chemiluminescent composition. In particular, the
present invention relates to improvements in such methods

which substantially increase light emission.

The invention further relates to the use of the chemiluminescent reactions and compositions in assay methods for detecting phosphatase enzymes and for detecting
5 phosphatase-labeled specific binding partners in immunoassays, nucleic acid probe assays and the like.

The present invention relates to a process for the preparation of chemiluminescent compounds which react with phosphatase enzymes to generate chemiluminescence. The
10 present invention relates to novel intermediates useful in this process. In particular, the present invention relates to a process and intermediates for preparing chemiluminescent compounds containing a heterocyclic ring group and an enol phosphate group which react with oxygen
15 upon removal of a phosphate protecting group to produce chemiluminescence and a carbonyl compound.

BACKGROUND OF THE INVENTION

a. Chemiluminescent Detection of Phosphatase Enzymes.

20 Hydrolytic enzymes such as alkaline phosphatase are frequently used as markers or labels in enzyme-linked assays for biological molecules and other analytes of interest such as drugs, hormones, steroids and cancer markers. In addition, phosphatase enzymes, e.g. alkaline
25 phosphatase (AP) and acid phosphatase (AcP), are clinically significant in their own right in human and veterinary diagnostics. Chemiluminescent detection of these enzymes offers a safe, convenient and sensitive means to provide a quantitative measure of the amount of enzyme in a sample or
30 of the amount of an enzyme-labeled analyte or labeled

specific binding partner for an analyte. Numerous chemiluminescent reaction schemes have been developed to quantitate the level of particular hydrolytic enzymes. Most of these schemes are complex and expensive, requiring
5 multiple enzymes or several reagents. Commercial acceptance of most of such methods for large volume testing has been slow.

Applicant's co-pending U.S. Patent Application Serial No. 08/585,090 which is fully incorporated herein by
10 reference, discloses the chemiluminescent reaction of certain heterocyclic compounds bearing an enol phosphate group with a phosphatase enzyme. Light emission is enhanced in the presence of cationic surfactants allowing the phosphatase to be detected at levels of 10^{-18} to 10^{-19} mol.

15 Applicant's co-pending U.S. Patent Application Serial No. 08/683,927 which is fully incorporated herein by reference, discloses the use of cationic aromatic compounds (CAC's) in conjunction with the chemiluminescent reaction of certain heterocyclic compounds bearing an enol phosphate
20 group with a phosphatase enzyme to substantially increase the amount of light emitted. The detection limit of phosphatase enzymes is thereby dramatically lowered.

b. Chemically and Enzymatically Triggerable Dioxetanes. Stable 1,2-dioxetanes bearing a protected
25 phenol group triggering group undergo a chemiluminescent decomposition upon removal of a protecting group (A. P. Schaap, T. S. Chen, R. S. Handley, R. DeSilva, and B. P. Giri, *Tetrahedron Lett.*, 1155 (1987); A. P. Schaap, R. S. Handley, and B. P. Giri, *Tetrahedron Lett.*, 935 (1987); A.
30 P. Schaap, M. D. Sandison, and R. S. Handley, *Tetrahedron*,

Lett., 1159 (1987); and A. P. Schaap, *Photochem. Photobiol.*, 47S, 50S (1988)). Enzymatically triggerable dioxetanes bear an aryloxy substituent which is blocked by an enzymatically removable protecting group. Reaction with a hydrolytic enzyme in an aqueous buffer reveals an aryloxy anion which accelerates the chemiluminescent decomposition rate of the dioxetane by orders of magnitude. Chemically triggerable dioxetanes bear an aryloxy substituent which is blocked by a protecting group which is removed by a simple chemical agent. An example is deprotection of an acetoxy dioxetane with hydroxide or a silyloxy dioxetane with fluoride. Numerous examples of such triggerable dioxetanes are disclosed, for example, in U.S. Patents 4,857,652, 5,068,339, 4,952,707, 5,112,960, 5,220,005, 5,326,882 and in PCT applications WO96/24849, WO94/10258 and WO94/26726. However, an inherent disadvantage of some triggerable dioxetanes is their tendency to generate background chemiluminescence in the absence of enzyme through slow thermal decomposition or non-enzymatic hydrolysis.

c. Luminol Derivatives. A phosphate and a NAG derivative of luminol are known (K. Sasamoto, Y. Ohkura, *Chem. Pharm. Bull.*, 38, 1323-5 (1991); M. Nakazono, H. Nohta, K. Sasamoto, Y. Ohkura, *Anal. Sci.*, 8, 779-83 (1992)). Treatment of the luminol derivative with the appropriate enzyme liberates luminol which is reacted in a subsequent step with ferricyanide to produce light.

d. Luciferin Derivatives. Phosphate and galactoside derivatives of firefly luciferin are known (N. Ugarova, Y. Vosny, G. Kutuzova, I. Dementieva, *Biolum. and Chemilum.*

New Perspectives, P. Stanley and L.J. Kricka, eds., Wiley, Chichester, 511-4 (1981); W. Miska, R. Geiger, J. Biolumin. Chemilumin., 4, 119-28 (1989)). Treatment of the firefly luciferin derivative with the appropriate enzyme liberates
5 firefly luciferin which is reacted in a second step with luciferase and ATP to produce light.

e. Reactions Involving the Generation of Reducing Agents. Chemiluminescent methods involving the generation of a reducing agent from a phosphate ester catalyzed by
10 alkaline phosphatase have been reported. (M. Maeda, A. Tsuji, K.H. Yang, S. Kamada, Biolum. and Chemilum. Current Status, 119-22 (1991); M. Kitamura, M. Maeda, A. Tsuji, J. Biolumin. Chemilumin., 10, 1-7 (1995); H. Sasamoto, M. Maeda, A. Tsuji, Anal. Chim. Acta, 306, 161-6 (1995)). The
15 reducing agent causes a reaction between oxygen and lucigenin to produce light arising from the lucigenin. Representative reducing agents include ascorbic acid, glycerol, NADH, dihydroxyacetone, cortisol and phenacyl alcohol. These methods are distinguished from the present
20 invention which involves the production of light from the deprotected fluorescent compound, not from lucigenin. The known methods of enzymatically generating a reducing agent for reaction with lucigenin all require a separate preliminary incubation step between the enzyme and the
25 phosphate compound. This adds additional complexity and assay time.

U.S. Patent 5,589,328 to Mahant discloses a chemiluminescent reaction whereby indoxyl esters, thioindoxyl esters and benzofuran esters are hydrolyzed by
30 an enzyme and thereby generate superoxide. Luminescence is

amplified by adding a chemiluminescence generating reagent such as lucigenin. Lucigenin produces chemiluminescence by reaction with superoxide.

f. Coupled Enzyme Methods. Numerous other

5 chemiluminescent methods and assays for determining hydrolytic enzymes such as phosphatase enzymes through coupled enzyme reactions are known. A compilation of such methods is listed in A. Tsuji, M. Maeda, H. Arakawa, Anal. Sci., 5, 497-506 (1989). Other examples of dual enzyme
10 chemiluminescent reactions are described in U.S. 5,306,621 and commonly assigned application Serial No. 08/300,367. The former describes the enzymatic generation of a peroxidase enhancer to enhance the chemiluminescent oxidation of luminol with a peroxidase; the latter
15 describes the enzymatic generation of a peroxidase enhancer to enhance the chemiluminescent oxidation of an acridancarboxylic acid derivative with a peroxidase.

With the exception of enzyme-triggered dioxetanes, each of the aforementioned methods suffers the drawback of
20 requiring multiple reagents or enzymes in order to generate the luminescent signal. The added expense or operational complexity has hindered commercial acceptance of these methods in spite of their demonstrated exceptional detection sensitivity. Chemiluminescent methods for
25 detecting and quantitating hydrolytic enzymes which achieve these levels of sensitivity but do not require additional enzymes or auxiliary reagents in addition to the enzyme substrate would be advantageous. The present invention provides such methods and compounds.

30

SUMMARY OF THE INVENTION


It is an object of the present invention to provide compounds for chemiluminescent detection of phosphatase enzymes which are thermally and hydrolytically stable at room temperature over an extended period of time and are
5 cleaved by a phosphatase enzyme to cleave the phosphate moiety.

It is also an object of the present invention to provide novel compounds substituted at one terminus of the
10 double bond with a nitrogen, oxygen or sulfur-containing heterocyclic ring group and further substituted at the other terminus of the double bond with an enzymatically cleavable phosphate (O-PO_3^{2-}) group which can be triggered to decompose with the generation of light.

15 It is a further object of the present invention to provide a method and compositions for generating chemiluminescence containing such novel compounds which can be triggered by a phosphatase enzyme.

It is yet another object of the present invention to
20 provide compounds which have superior light-generating ability and provide significant advantages when used for the detection of phosphatase enzymes, and for use in immunoassays and the detection of enzyme-linked nucleic acids, antibodies, haptens and antigens by generally known
25 methods which employ phosphatase labels for detection of analytes.

The above and other objects and advantages in accordance with the present invention are attained by a compound having the formula I:



5 wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from N, O and S atoms, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group and wherein each M is
10 independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality.

The above and other objects and advantages in accordance with the present invention are further attained by a reagent composition which produces chemiluminescence
15 in the presence of a phosphatase enzyme which comprises in an aqueous solution: a compound of formula I and at least one surfactant enhancer in an amount effective to enhance the chemiluminescence.

The above and other objects and advantages in accordance with the present invention are further attained
20 by a method for producing chemiluminescence which comprises reacting a phosphatase enzyme with at least one compound of formula I.

The above and other objects and advantages in accordance with the present invention are further attained by a method
25 for detecting an analyte in a sample by a chemiluminescent assay procedure which comprises: reacting a phosphatase enzyme with at least one compound of formula I to produce chemiluminescence for detecting the analyte; detecting the

chemiluminescence; and relating the amount of the chemiluminescence to the amount of the analyte.

The above and other objects and advantages in accordance with the present invention are further attained by a method
5 of detecting an analyte in an assay procedure by a chemiluminescent reaction which comprises: providing a reagent composition which generates chemiluminescence in the presence of a phosphatase enzyme which comprises, in an aqueous solution, at least one compound of formula I which
10 reacts with the phosphatase enzyme wherein in the compound of formula I, and a surfactant enhancer in an amount effective to enhance the chemiluminescence; reacting a phosphatase enzyme with the composition to produce chemiluminescence for detecting the analyte; and relating
15 the amount of chemiluminescence to the amount of the analyte.

It is also an object of the present invention to provide compositions comprising a cationic aromatic compound and a compound of formula I which can be triggered to decompose
20 with the generation of light.

It is a further object of the present invention to provide an improved method for generating chemiluminescence by reaction of a reagent composition with a phosphatase enzyme.

25 Still further, it is an object of the present invention to provide a composition and method which rapidly produces efficient chemiluminescence on reaction with a phosphatase enzyme.

The above and other objects and advantages in accordance
30 with the present invention are attained by a reagent

composition comprising a cationic aromatic compound and a compound of formula I.

The above and other objects and advantages in accordance with the present invention are further attained by a reagent composition which produces chemiluminescence in the presence of a phosphatase enzyme which comprises in an aqueous solution: a cationic aromatic compound, a compound of formula I which reacts with the phosphatase enzyme and, in combination, at least one anionic surfactant and at least one non-ionic surfactant in amounts effective to provide rapid generation of efficient chemiluminescence.

It is a further object of the present invention to provide a synthetic process and intermediates useful therein for the preparation of chemiluminescent compounds which react with phosphatase enzymes to generate light.

It is a further object of the present invention to provide a process and intermediates for preparing chemiluminescent compounds of formula I containing a heterocyclic ring group and an enol phosphate group which reacts with oxygen upon removal of a phosphate protecting group to produce light and a carbonyl compound.

It is a further object of the present invention to provide a process comprising phosphorylating an enolate of an ester or thioester compound to produce a phosphodiester or triester intermediate compound which is deprotected to a phosphate monoester salt.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph showing the time profile of the chemiluminescence intensity emitted by 100 μ L of a reagent

consisting of a 0.33 mM solution of acridan phosphate **1**
(9-(phenoxyphosphoryloxymethylidene)-10-methylacridan,
disodium salt) in 0.1 M tris buffer, pH 8.5 and 0.01 mg/mL
of the enhancer polyvinylbenzyltributylphosphonium chloride
5 co-polyvinylbenzyltrioctylphosphonium chloride (containing
about a 3:1 ratio of tributyl: trioctyl groups), (Enhancer
A) triggered at 25 °C by addition of 8×10^{-16} mol of
alkaline phosphatase (AP). The figure also shows for
comparison the chemiluminescence profile of the ester
10 phenyl 10-methylacridan-9-carboxylate in a similar reagent
composition containing the ester in place of **1**.

Figure 2 is a plot of the ultraviolet-visible absorp-
tion spectrum of a reaction mixture containing 3 mL of a
15 reagent composition containing **1** with 2.4×10^{-12} mol of AP
at room temperature. Curves labeled A-F represent scans
taken 0, 5, 10, 20, 40 and 60 min after addition of enzyme.

Figure 3 is a plot of the chemiluminescence spectrum
20 produced by reaction of 200 μ L of a reagent composition
containing **1** with 8×10^{-13} mol of AP at room temperature.

Figure 4 is a graph relating the amount of AP to the
maximum chemiluminescence intensity emitted by 100 μ L of a
25 reagent containing acridan phosphate **1** triggered at room
temperature. Chemiluminescence emission was initiated at
room temperature by addition of 10 μ L of solutions of AP
containing between 8×10^{-15} mol and 8×10^{-20} of enzyme to
100 μ L of a 0.33 mM solution of acridan phosphate **1** in tris
30 buffer, 0.1 M (pH 8.5) containing 0.88 mM Mg^{+2} and 0.01

mg/mL of Enhancer A in the wells of a white microplate. The term S-B refers to the chemiluminescence signal (S) in Relative Light Units (RLU) in the presence of AP corrected for background chemiluminescence (B) in the absence of AP.

- 5 The graph shows linear detection of alkaline phosphatase. The calculated detection limit (twice the standard deviation of the background) was determined to be 8×10^{-19} mol under these conditions.

- 10 Figure 5 is a graph relating the amount of AP to the total chemiluminescence intensity emitted by 100 μ L of a reagent composition comprising 0.33 mM acridan phosphate 1 in 0.1 M tris buffer, pH 9, 0.88 mM $MgCl_2$. The composition was reacted with 10 μ L of solutions of AP containing
- 15 between 8×10^{-15} and 8×10^{-20} mol of enzyme or 10 μ L of water as a reagent blank for 4 min at ambient temperature in the wells of a black microplate. A solution containing 0.5 mg/mL of Enhancer A in 1 N NaOH (100 μ L) was injected and light intensity integrated for 10 sec. The graph shows
- 20 linear detection of alkaline phosphatase.

- Figure 6 is a digitally scanned image of an x-ray film from a Western blot experiment detecting human transferrin via an AP-labeled antibody on a PVDF membrane with a
- 25 chemiluminescent reagent composition. Dilutions of transferrin containing from 5000, 1000, 180, 30 and 5 pg, respectively, of protein were electrophoresed and transferred to the membrane. Protein was detected by briefly soaking the blot with a reagent composition containing 0.5
- 30 mg/mL of Enhancer A in 0.2 M 2-amino-2-methyl-1-propanol,

buffer, pH 9.6 containing 0.88 mM MgCl_2 and 0.66 mM acridan phosphate 1 and exposing to x-ray film for 1 min after a 10 min incubation time.

5 Figure 7 is a graph showing the effect on chemiluminescence production from reaction of 5 with alkaline phosphatase (AP) by adding lucigenin to the reaction solution. Equal volumes of two solutions containing 5 in buffer were reacted with 8×10^{-16} moles of alkaline phosphatase (AP). One solution also contained 6.4 μM lucigenin.

15 Figure 8 is a graph showing the time profile of the chemiluminescence intensity emitted by 100 μL of Reagent A consisting of 0.1 M tris buffer, pH 8.8, 6.4 μM lucigenin, 0.66 mM acridan phosphate 5 (from a 1:100 dilution of a 0.066 M methanol solution), 1 mg/mL sodium dodecyl sulfate, 0.01 mg/mL Na_2SO_3 , 0.033% (w/v) TWEEN 20 and 0.88 mM MgCl_2 triggered at room temperature by addition of 8×10^{-16} mol of AP. The figure also shows for comparison the chemiluminescence profile from a reagent composition containing 5 and a cationic polymeric surfactant enhancer (0.25TB/TO).

25 Figure 9 is a graph relating the amount of AP to the maximum chemiluminescence intensity emitted by 100 μL of a reagent composition of the present invention (reagent A) containing acridan phosphate 5, lucigenin, SDS and TWEEN 20 triggered at 25 $^\circ\text{C}$. For comparison, a similar graph is presented from a set of experiments using a reagent composition (reagent B) containing the same acridan phosphate and (0.25TB/TO) but without lucigenin triggered at 37 $^\circ\text{C}$.

Chemiluminescence emission was initiated by addition of 10 μL of solutions of AP containing between 8×10^{-15} and 8×10^{-22} mol of enzyme to 100 μL of the respective reagent compositions in the wells of a white microplate. The term
5 S-B refers to the chemiluminescence signal (S) in Relative Light Units (RLU) in the presence of AP corrected for background chemiluminescence (B) in the absence of AP. The graph shows that reagent A including lucigenin achieves a 100-fold lower detection limit for AP.

10

Figure 10 is a graph relating the amount of AP to the maximum chemiluminescence intensity emitted by 100 μL of a reagent composition comprising 0.66 mM acridan phosphate 7 in 0.1 M tris buffer, pH 8.8, 6.4 μM lucigenin, 1 mg/mL
15 sodium dodecyl sulfate, 0.01 mg/mL Na_2SO_3 , 0.033% (w/v) TWEEN 20 and 0.88 mM MgCl_2 . The composition was reacted with 10 μL of solutions of AP containing between 8×10^{-15} and 8×10^{-22} mol of enzyme or 10 μL of water as a reagent blank and measured after 75 sec at ambient temperature in
20 the wells of a black microplate.

Figure 11 is a graph relating the amount of AP to the maximum chemiluminescence intensity emitted by 100 μL of a reagent composition comprising 0.66 mM acridan phosphate 8
25 in 0.1 M tris buffer, pH 8.8, 6.4 μM lucigenin, 1 mg/mL sodium dodecyl sulfate, 0.01 mg/mL Na_2SO_3 , 0.033% (w/v) TWEEN 20 and 0.88 mM MgCl_2 . The composition was reacted with 10 μL of solutions of AP containing between 8×10^{-15} and 8×10^{-22} mol of enzyme or 10 μL of water as a reagent
30 blank and measured after 75 sec at ambient temperature in

the wells of a black microplate.

Figure 12 is a graph showing the rapid generation of chemiluminescence emitted by 100 μ L of a reagent consisting of 0.1 M tris buffer, pH 8.8, 6.4 μ M lucigenin, 0.66 mM acridan phosphate [(from a 1:100 dilution of a 0.066 M methanol solution), 1 mg/mL sodium dodecyl sulfate, 0.01 mg/mL Na_2SO_3 , 0.033% (w/v) TWEEN 20 and 0.88 mM MgCl_2 triggered at room temperature by addition of 8×10^{-16} mol of AP.

Figure 13 is a graph showing the results of a chemiluminescent immunoassay for human chorionic gonadotropin using a detection reagent of the invention.

Figure 14 is a graph showing the results of a rapid chemiluminescent immunoassay for thyroid stimulating hormone using a detection reagent of the invention.

Figure 15 is a graph showing the results of a chemiluminescent immunoassay for thyroid stimulating hormone using a detection reagent of the invention.

Figure 16 is a graph showing the results of a chemiluminescent immunoassay for estradiol using a detection reagent of the invention.

Figure 17 is a plot demonstrating the ability to simultaneously detect acid phosphatase and alkaline phosphatase in one sample. The upper trace is the light emitted

from a sample containing both enzymes, the lower trace is the light emitted from a sample containing only acid phosphatase and the middle trace represents the difference attributable to alkaline phosphatase.

5


Figure 18 is a graph relating the amount of AP to the maximum chemiluminescence intensity emitted by 100 μ L of a reagent composition comprising 0.66 mM acridan phosphate 12 in 0.1 M tris buffer, pH 8.8, 6.4 μ M lucigenin, 1 mg/mL sodium dodecyl sulfate, 0.01 mg/mL Na_2SO_3 , 0.033% (w/v) 10 TWEEN 20 and 10 μ M MgCl_2 . The composition was reacted with 10 μ L of solutions of AP containing between 8×10^{-15} and 8×10^{-22} mol of enzyme or 10 μ L of water as a reagent blank and measured after 75 sec at ambient temperature in the 15 wells of a black microplate.

Figure 19 is a graph relating the amount of AP to the maximum chemiluminescence intensity emitted by 100 μ L of a reagent composition comprising 0.33 mM acridan phosphate 13 20 in 0.1 M tris buffer, pH 8.8, 3.2 μ M lucigenin, 0.5 mg/mL sodium dodecyl sulfate, 0.005 mg/mL Na_2SO_3 , 0.016% (w/v) TWEEN 20 and 5 μ M MgCl_2 . The composition was reacted with 10 μ L of solutions of AP containing between 8×10^{-15} and 8×10^{-22} mol of enzyme or 10 μ L of water as a reagent blank 25 and measured after 75 sec at ambient temperature in the wells of a black microplate.


DESCRIPTION OF THE PREFERRED EMBODIMENTS

It has been unexpectedly discovered as disclosed in Applicant's co-pending U.S. Patent Application Serial No.

08/585,090 that certain novel compounds react with a phosphatase enzyme to generate easily detectable chemiluminescence. Compounds of the present invention which produce chemiluminescence in the presence of a phosphatase enzyme have the formula:





wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from N, O and S atoms, wherein Z is an O or S atom, wherein R_6 is an organic group which allows the light to be produced and wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality. Reaction of the phosphatase with I in the presence of oxygen leads to formation of a carbonyl-containing compound VI in an electronically excited state (VI^*) and the dephosphorylated product VII. Radiative decay of VI^* leads to light emission.



The heterocyclic ring system of I contains at least one heteroatom selected from N, O and S atoms and is in

conjugation with the ring carbon bearing the exocyclic double bond. Preferred heterocyclic compounds include compounds of formulas II and III as depicted below and their double bond isomers or mixtures of the isomers.








19


where Z, M, n and R₆ are as defined above and R₁ - R₅ are defined below.

Referring back to formula I, exemplary ring structures which can comprise the group Het include the structures below where the asterisk denotes the position of the exocyclic double bond. Without explicitly showing all possible substitution patterns, it is to be understood that each ring position can contain substituents other than hydrogen. Other heterocyclic ring compounds not specifically listed below but still falling within the scope of formula I will occur to the skilled artisan.

25







5

In all of the above compounds, the group R_1 is an organic group containing from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms exclusive of the necessary number of H atoms required satisfy the valencies of the atoms in the group. The organic group is preferably selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups. More preferred groups for R_1 include $C_1 - C_4$ alkyl groups and benzyl groups.

In all of the above compounds, the groups $R_2 - R_5$, which can be the same or different, each are H or a substituent group which permits the light to be produced and will generally contain from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms. Representative substituent groups which can be present include, without limitation, alkyl, substituted alkyl, aryl, substituted aryl, aralkyl, alkenyl, alkynyl, alkoxy, aryloxy, halogen, amino, substituted amino groups, carboxyl, carboalkoxy, carboxamide, cyano, and sulfonate groups. Either or both pairs of adjacent groups, i.e. $R_2 - R_3$ or $R_4 - R_5$, can be joined together as a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring which

is fused to the ring bearing the exocyclic double bond. Such fused heterocyclic rings can contain 1 or more N, O or S atoms and can be substituted at the ring carbons with groups other than hydrogen such as those mentioned above.

5 Substituent groups can be incorporated in various quantities and at selected ring positions in order to modify the properties of the compound or to provide for convenience of synthesis of the final phosphate compound. Such properties include, for example, chemiluminescence
10 quantum yield, rate of reaction with the enzyme, maximum intensity of light emission, duration of light emission, wavelength of light emission and solubility in the reaction medium. Specific substituents and their effects are illustrated in the specific examples below, which,
15 however, are not to be considered limiting the scope of the invention in any way.

Each of the groups M independently comprise a hydrogen atom or a cationic center. A cationic center means a positively charged atom such as a sodium atom Na^+ , a group
20 of atoms such as an ammonium ion NH_4^+ or a portion of a molecule with one or more sites of positive charge. Examples of the latter include dicationic compounds described in U.S. Patent 5,451,347 to applicant and polymeric compounds with multiple cationic groups as
25 described in applicant's U.S. Patent 5,393,469. the positive charge on a cationic center may take any unit value, i.e. 1, 2, 3 etc. Exemplary cationic centers include, without limitation, alkali metal ions, alkaline


earth ions, quaternary ammonium ions and quaternary phosphonium ions and are present in the number required by their valence. If two groups are required to be present in the compound of formula I for electroneutrality, they can
5 be the same or different. Preferred counter ions are the alkali metal ions.

As noted above, the organic group R_6 can be any group which allows or does not interfere with the light production and preferably contains from 1 to 50 atoms
10 selected from C, N, O, S, P and halogen atoms and the necessary number of H atoms required satisfy the valencies of the atoms in the group. Groups which can function as the R_6 group include, without limitation, alkyl, substituted alkyl, aryl, substituted aryl and aralkyl
15 groups. Substituent groups other than H atoms, such as ionic groups or polar groups, can be incorporated in various numbers and at selected positions on the carbon chain or ring of R_6 in order to modify the properties of the compound or to provide for convenience of synthesis of
20 the final phosphate compound. Such properties include, for example, chemiluminescence quantum yield, rate of reaction with the enzyme, maximum intensity of light emission, duration of light emission, wavelength of light emission and solubility in the reaction medium. Specific
25 substituents and their effects are illustrated in the specific examples below, which, however, are not to be considered limiting the scope of the invention in any way.

A preferred class of compounds have the formula IV

below wherein Ar is an aryl ring group containing at least one carbocyclic or heterocyclic aromatic ring and which can be further substituted, Z is selected from O and S atoms, R₁ is as defined above, M is H or a monovalent or divalent cationic counter ion as defined above and n is 2 or 1, respectively.


10



The groups R₇ to R₁₄, which can be the same or different, each are H or a substituent group containing from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms and which permit the light to be produced and can include, without limitation, alkyl, substituted alkyl, aryl, substituted aryl, aralkyl, alkenyl, alkynyl, alkoxy, aryloxy, halogen, amino, substituted amino groups, carboxyl, carboalkoxy, carboxamide, cyano, and sulfonate groups. Adjacent groups, e.g. R₇ - R₈ or R₈ - R₉, can be joined together as a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring. It is preferred that R₇ to R₁₄ are selected from hydrogen and alkoxy groups such as methoxy, ethoxy, t-butoxy and the like. Especially preferred compounds have the formula V where Z is O or S and Ar is a phenyl group, a substituted phenyl group, a naphthyl group or a substituted naphthyl

group and M and n are as defined above.


5



V

Another aspect of the present invention is the use of compounds of any of formulas I-V in a method to produce visible chemiluminescence by reaction with a phosphatase enzyme. Reaction of a compound of formula I-V with a phosphatase enzyme in an aqueous buffer solution produces easily detected chemiluminescence. Light intensity reaches a maximum level within minutes at room temperature when the reaction is conducted at alkaline pH. The reaction is conducted optionally in the presence of an enhancer.

20




While not wishing to be bound by any specific mechanistic explanation for this discovery at this point, molecular oxygen is a necessary reactant and the reaction products are ketone VI, an ester or thioester VII, a compound of the formula R_6ZH and phosphate ion. Light is emitted from the electronically excited state of VI. A necessary condition for the production of light is that the reaction produces sufficient energy to form the

excited state of VI. If VI is fluorescent then chemiluminescence is produced directly from the reaction via emission from the excited state of VI. An especially surprising finding is that reaction of compounds of

5 formula I with a phosphatase enzyme at moderately basic pH produces far more intense light emission than is produced by autoxidation of the enolate intermediate (the anion of VII) which would be expected to form upon cleavage of the phosphate group.

10 In a preferred method of producing chemiluminescence, a compound containing an acridan ring is reacted with alkaline phosphatase in an alkaline buffer with a pH between about 8 and 10 to produce a continuous chemiluminescence signal which commences upon reaction of

15 the enzyme and the phosphate compound. The light intensity at any time point can be increased up to at least 40-fold by incorporation of at least one surfactant enhancer as will be described in more detail below.



In a preferred method of producing light from the reaction of compound IV with a phosphatase enzyme, the reaction is performed at a temperature between 5 °C and 50

°C, preferably between 20 °C and 40 °C in an aqueous buffer solution at a pH between 7 and 10.5, preferably between 8.5 and 10. Compound IV is used at a concentration between 1 μ M and 20 mM, preferably between 10 μ M and 1 mM.

- 5 The enzyme is preferably an alkaline phosphatase or an alkaline phosphatase conjugate. Light is emitted from the excited state of VIII.

Compounds of the present invention typically produce light over a 100-200 nm wide band of emission, which
10 exhibits a maximum intensity at wavelengths in the near ultraviolet to the visible region of the electromagnetic spectrum. Typical wavelengths of maximum intensity λ_{max} in the range of 350-500 nm. It is contemplated that phosphate compounds of formula I bearing a covalently linked
15 fluorophore not in conjugation with the double bond of the vinyl phosphate moiety could, upon formation of the excited product VI*, undergo intramolecular energy transfer resulting in emission at longer wavelengths from the excited state of the fluorophore.

- 20 More than one compound of formula I can be used concurrently in a method for producing light by the action of a phosphatase enzyme. It may be advantageous in some instances to simultaneously react two or more compounds of formula I with the phosphatase enzyme. When the two or
25 more compounds have differing luminescent or physical properties, the combination of the two may be desirable to produce a light emitting reaction with characteristics not readily achievable through the use of any one compound.

Examples of luminescent and physical properties which can differ between compounds I include emission spectrum, duration of light emission, enzyme turnover, rate of rise of emission to maximum, hydrophobicity/hydrophilicity and solubility. While particular luminescent properties can differ among the compounds of formula I in the present methods, the variation in properties does not detract from the basic utility of the compounds; selection of particular compounds with desirable properties can be made by virtue of the teachings and methods described herein.

Light emitted by the present method can be detected by any suitable known means such as a luminometer, x-ray film, high speed photographic film, a CCD camera, a scintillation counter, a chemical actinometer or visually. Each detection means has a different spectral sensitivity. The human eye is optimally sensitive to green light, CCD cameras display maximum sensitivity to red light, x-ray films with maximum response to either UV to blue light or green light are available. Choice of the detection device will be governed by the application and considerations of cost, convenience, and whether creation of a permanent record is required.

It is contemplated that fluorescent energy acceptors can be employed to shift the maximum emission to longer wavelengths (red-shifting). Various techniques for red-shifting emission are known in the art of chemiluminescent reactions and assays. Covalently linked fluorophores as described above are one example. Fluorescers can be added

to the reaction solution as separate species. Fluorescers can be linked to an enhancer substance such as a cationic polymer or associated with an enhancer substance such as a micelle or polymer in order to bring the fluorescer in
5 close contact to the compound. Alternately, the fluorescer can be provided in a non-fluorescent form which is convertible to the fluorescent form by removal of a phosphate group during the enzyme reaction period. Examples of the latter type of compound include
10 fluorescein diphosphate, coumarin phosphates such as 4-methylumbelliferone phosphate, benzothiazole phosphates such as ATTOPHOS (JBL Scientific, San Luis Obispo, California).

At least one enhancer compound can also be employed in
15 the chemiluminescent reaction of the present invention, in order to increase the amount of light emitted. Enhancers which are effective in the present method can function by increasing the fraction of excited state product molecules which emit light, by increasing the fraction of product
20 molecules which are formed in the excited state, by increasing the rate of reaction or turnover of the enzyme, by increasing the rate of a subsequent chemical reaction step, by improving the stability of the enzyme, by promoting the association of the enzyme with the compound
25 of formula I, by inhibiting or preventing competitive non-luminescent side reactions or by any combination of these mechanisms. Enhancers will be used in an amount effective to enhance the chemiluminescence, preferably between 0.001

and 5 mg/mL in the final reaction solution, more preferably between 0.01 and 2.5 mg/mL.

Enhancer compounds that are effective in the practice of the present method are typically surfactant compounds, i.e. compounds which display surface active properties, e.g., surface tension depression, surface wetting or detergency. Surfactants comprise a hydrophilic region containing polar and/or ionic groups and a hydrophobic region containing mainly hydrocarbon groups or alkylenoxy groups or both. Surfactants are categorized as cationic, anionic, nonionic and zwitterionic and can be monomeric or polymeric. Cationic surfactant enhancers found useful in the practice of the present invention include polyvinyl type polymers with pendant quaternary phosphonium groups which are disclosed in U.S. Patent 5,393,469 the disclosure of which is incorporated herein by reference. Exemplary polymers of this type include polyvinylbenzyltributylphosphonium chloride copolymer with polyvinylbenzyltrioctylphosphonium chloride and (polyvinylbenzyltributylphosphonium chloride). Polyvinyl type polymers with pendant quaternary ammonium groups are also useful as enhancers in the present invention. Examples of such polymers are disclosed in U.S. Patent 5,112,960, the disclosure of which is incorporated herein by reference and include polyvinylbenzylbenzyltrimethylammonium chloride and polyvinylbenzyltributylammonium chloride.

Another category of cationic surfactant enhancers

found useful in the practice of the present invention are dicationic compounds bearing two quaternary ammonium or phosphonium groups as disclosed in U.S. Patent 5,451,347 the disclosure of which is incorporated herein by

5 reference. The compound (1-trioctylphosphoniummethyl-4-tributylphosphoniummethylbenzene dichloride) for example provides enhanced chemiluminescence when used in a method for producing light according to the present invention. Still other cationic surfactant enhancers useful in the
10 practice of the present invention are monoquaternary ammonium salts e.g. cetyltrimethylammonium chloride or bromide and monoquaternary phosphonium salts such as cetyltrimethylammonium bromide.

Anionic surfactant enhancers include alkyl sulfates
15 and alkylsulfonates. Nonionic surfactant enhancers include polyoxyethylenated alkylphenols, polyoxyethylenated alcohols, polyoxyethylenated ethers and polyoxyethylenated sorbitol esters. Zwitterionic surfactant enhancers include quaternary ammoniumalkyl phosphates and sulfonates. More
20 extensive lists of exemplary structures of each category of surfactant can be found in any standard treatise on surfactants. Numerous representative member surfactants have been tested and found to be effective to varying degrees in increasing the amount or intensity of light
25 produced compared to the amount produced in its absence. Cationic surfactants, in particular, quaternary ammonium and quaternary phosphonium salts have been found to be among the most effective enhancers.

As described above, surfactant enhancers can also have fluorescent groups covalently attached or associated through electrostatic or hydrophobic interactions.


In an alternate mode of performing the present
5 chemiluminescent reactions, the chemiluminescent compound is reacted with a phosphatase enzyme in a buffer at a first pH in the range 5.0 - 9.5 for a first period of time ranging from a few seconds to less than about 10 min in the absence of an enhancer. Any light produced during this
10 first period is neglected. Then a strongly basic trigger solution containing an enhancer is added at once and the burst of light measured either by measuring the peak intensity, or integrating for a second fixed time period or until light emission has ceased. It is desirable that
15 the pH and amount of enhancer be selected so that all of the light is emitted in a short period of time, preferably about one minute or less. The pH of the trigger solution should be > 11 and preferably above 12. Preferred bases are sodium hydroxide and potassium hydroxide. Enhancers
20 useful in this mode of reaction are as identified in the foregoing discussion and will be used in an amount effective to enhance the chemiluminescence, preferably between 0.001 and 1 mg/mL in the final reaction solution.

This mode of generating the chemiluminescence may be
25 advantageous in assays in which a large number of samples is processed simultaneously for measurement at a later time. Detection of acid phosphatase can also be performed in this manner. An optional step which can also be

incorporated into such a chemiluminescent reaction or assay is to add an enzyme inhibitor to the reaction system after the first period of time to stop all further enzyme action.

5 Since the reaction is catalyzed by the phosphatase enzyme, exceedingly small quantities of the enzyme are sufficient to produce a detectable amount of light. Sensitivities of 1 attomol (1×10^{-18} mol) have been achieved. The ability to detect such small amounts of
10 phosphatase enzymes make the present chemiluminescent technology suitable for analyses of many types of analytes using enzyme-linked assays.

 The chemiluminescent reaction of the present invention provides a particularly effective reagent for detection of
15 AP conjugates on polyvinylidene difluoride (PVDF) membranes, nylon membranes and on nitrocellulose filters and membranes. Surprisingly, it has been found that reaction of an antibody-AP conjugate on a PVDF membrane with a reagent composition containing compound 1



1

20 provides an intense luminescence which rises to a maximum
25 level nearly instantly and maintains essentially constant intensity for at least 2 days. Detection on film and

optimizing image intensity are thus particularly convenient.

Surprisingly, it has been found that light emission produced on PVDF membranes persists at useful levels for very long periods of time, exceeding a month. Art-known chemiluminescent reagents used in membrane-based detection schemes provide a luminescent signal with a time course which shows either a rapid decay, e.g. luminol with peroxidase which decays within 3-4 hours, or a slow rise and gradual decay as is the case with enzyme-triggered dioxetanes.

In another aspect, the present invention relates to a reagent composition for producing chemiluminescence by reaction with a phosphatase enzyme comprising an aqueous buffer with a pH between about 7 and about 10.5, a compound of formula I at a concentration of 0.01-10 mM and optionally at least one enhancer in an amount effective to enhance the chemiluminescence, preferably between 0.001 and 10 mg/mL. Formulations for chemiluminescent reaction with alkaline phosphatase can further comprise a magnesium or zinc salt at a concentration of 0.01-10 mM for increasing the activity of the enzyme.


A preferred reagent composition for producing chemiluminescence by reaction with a phosphatase enzyme comprises an aqueous buffer with a pH between about 7 and about 10.5, acridan phosphate of formula IV or V at a concentration of 0.01-10 mM and a surfactant enhancer in an amount effective to enhance the chemiluminescence,

preferably between 0.001 and 10 mg/mL. The formulation further comprises a magnesium salt at a concentration of 0.01-10 mM.

It is preferred that the surfactant enhancer in the reagent composition is selected from the group consisting of polymeric cationic enhancers containing ammonium or phosphonium groups and dicationic enhancers containing ammonium or phosphonium groups. Especially preferred are polyvinyl polymers bearing a pendant trialkylphosphonium group on each monomeric unit and polyvinyl polymers bearing a pendant trialkylammonium group or a benzyldialkylammonium group on each monomeric unit. The amount and choice of enhancer can be selected for optimum performance in a given application as a matter of routine experimentation. A preferred composition for detection of AP or conjugates in solution comprises an amine buffer, pH 8.5 the compound 9-(phenoxyphosphoryloxymethylidene)-10-methylacridan, disodium salt, 0.1-1 mM, a magnesium salt, 0.1-1 mM, and Enhancer A (polyvinylbenzyltributylphosphonium chloride co-polyvinylbenzyltrioctylphosphonium chloride containing about a 3:1 ratio of tributyl:trioctyl groups), 0.01-0.1 mM. A preferred composition for detection of AP or conjugates on a membrane comprises an amine buffer, pH 9.6, the compound 9-(phenoxyphosphoryloxymethylidene)-10-methylacridan, disodium salt, 0.1-1 mM, a magnesium salt, 0.1-1 mM, and Enhancer A, 0.1-1 mM.

Applicant's co-pending U.S. Patent Application Serial No. 08/683,927 discloses that adding a cationic aromatic compound (CAC) to the above reaction system greatly increases the quantity and/or intensity of light produced.

5 Further, adding the CAC to the compound of formula I in the absence of the phosphatase enzyme does not lead to a corresponding increase in spontaneous (background) chemiluminescence since it is believed that the CAC exerts its effect on a dephosphorylated intermediate derived from
10 compound I and not on compound I itself. Greatly increased sensitivity of detection of the phosphatase enzyme results by virtue of the increased signal/background. It is significant that light emission continues to arise from the excited state of VI and not from the CAC.




20 A variety of CACs have been found to function to increase the quantity and/or intensity of light produced. CACs are aromatic compounds bearing at least one positive charge either on the aromatic ring or ring system or residing on a substituent on one of the rings, provided
25 that the substituent is in conjugation with the unsaturated ring atoms.

The CACs of the present invention are those compounds having oxidation/reduction potentials suitable for causing

an increase in chemiluminescence in reactions of the invention. The suitability of compounds as CACs can be readily determined by means of the methods set forth in the specific examples below. Without being bound by any particular mechanistic interpretation, it appears that an intermediate product of dephosphorylation of I undergoes a redox reaction which can be reversible or irreversible. Molecular oxygen reacts with one or more of the reacting species in the reaction selected from the CAC, a reduced form of the CAC or the dephosphorylated intermediate of compound I or an oxidized or reduced form thereof to ultimately form an oxygenated reaction product derived from compound I. The oxygenated reaction product undergoes a chemiluminescent reaction, which is likely an O-O bond breaking reaction.

The CACs of the present invention can be a heteroaromatic ring compound comprising one or more isolated or fused aromatic rings containing at least one atom other than carbon (heteroatom), preferably one or more nitrogen atoms in which the positive charge is substantially localized on one or more of the heteroatoms. Examples of this class of CAC are cyanine dyes, thiacyanine dyes, carbocyanine dyes, thiacyanocyanine dyes, selenacyanocyanine dyes, azo dyes and acridinium derivatives of the formula:



5 wherein Q is an electron withdrawing group, X⁻ is a non-interfering anionic counter ion and R is an alkyl or aralkyl group each of which can optionally contain non-interfering substituents. Acridinium derivatives in which some of the ring hydrogens are replaced by other

10 substituting groups are also within the scope of functional CACs. The electron withdrawing group Q can be e.g. a halogen atom such as Cl, a cyano group, or a carbonyl group such as an ester group -COOR, a thioester group -COSR, an amide group -CONR¹R² or a sulfonimide group

15 -CON(R)SO₂R'. Similarly the CAC can be a derivative of a phenanthridinium or phenanthroline compound.

The CACs of the present invention can be an aromatic ring compound comprising one or more isolated or fused carbocyclic aromatic rings bearing at least one cationic

20 substituent containing at least one heteroatom, provided that the cationic site is in conjugation with the aromatic ring. In the latter class of CAC it is preferred that the heteroatom be a nitrogen or sulfur atom. Examples include Methylene Blue and Nile Blue.

25 The CACs of the present invention can bear more than one positive charge; e.g. dicationic compounds are within the scope of the invention. Exemplary compounds of this type include N,N'-dimethylbiacridinium dinitrate, commonly

known as lucigenin and 1,1'-dimethyl-4,4'-bipyridinium dichloride, commonly known as methyl viologen dichloride, or paraquat dichloride.


Additional compounds which are useful as the CAC component of the present invention include, by way of illustration, compounds in Table I.

Table 1. CACs

- Alcian Yellow, Basic Blue 41, Basic Blue 66, Basic Red 29, 3-Benzyl-5-(2-hydroxyethyl)-4-methylthiazolium chloride,
- 10 [2-[2-[2-Chloro-3-[(1,3-dihydro-1,3,3-trimethyl-2H-indol-2-ylidene)ethylidene]-1-cyclohexen-1-yl]ethenyl]1,3,3-trimethylindolium perchlorate, {IR-786 perchlorate}, trans-4-[4-(Dibutylamino)styryl]-1-methylpyridinium iodide,
- 15 5,5'-Dichloro-11-diphenylamino-3,3'-diethyl-10,12-ethylene thiatricarbocyanine perchlorate, {IR-140}, 3,3'-Diethyl-9-methylthiacarbocyanine iodide, 1,1'-Diethyl-2,2'-quinotricarbocyanine iodide, 3,3'-Diethylselenacarbocyanine iodide,
- 20 3,3'-Diethylthiacyanine iodide, 3,3'-Diethylthiadibocarbocyanine iodide, 2-[4-(Dimethylamino)styryl]-3-ethylbenzothiazolium iodide, 3,6-Dimethyl-2-(4-dimethylaminophenyl)-benzothiazolium bromide,
- 25 3,4-Dimethyl-5-(2-hydroxyethyl)thiazolium iodide, 4-[2-[3-[(2,6-Diphenyl-4H-thiopyran-4-ylidene)ethylidene]-2-phenyl-1-cyclo-hexen-1-yl]ethenyl]-2,6-diphenylthiopyrylium tetrafluoroborate {IR-1040},


- 5-[3-Ethoxy-4-(3-ethyl-5-methyl-2(3H)-benzothiazolyl-
idene)-2-butenylidene]-3-ethyl-2-[(3-ethyl-4,5-
diphenyl-2(3H)-thiazolylidene)methyl]-4,5-dihydro-4-
oxothiazolium iodide,
- 5 3-Ethyl-2-(2-hydroxy-1-propenyl)benzothiazolium chloride,
3-Ethyl-2-methylbenzothiazolium iodide,
3-Ethyl-2-methylbenzoxazolium iodide,
1-Ethyl-3-methyl-1H-imidazolium chloride,
Methylene Blue, Nile Blue A, and
- 10 Triphosphopyridine nucleotide, sodium salt hydrate

Another aspect of the present invention, therefore, is
the use of compounds of any of formulas I-V and a CAC in a
method to produce detectable chemiluminescence by reaction
15 with a phosphatase enzyme. Reaction of a compound of
formula I-V with a phosphatase enzyme in the presence of a
CAC in an aqueous buffer solution produces easily detected
chemiluminescence from the excited state of VI. Light
intensity reaches a maximum level within seconds to
20 minutes at room temperature when the reaction is conducted
at alkaline pH.



In a preferred method of producing chemiluminescence,
a compound of formula IV containing an acridan ring is

reacted in the presence of a CAC with alkaline phosphatase in an alkaline buffer with a pH between about 7 and 10.5 to produce a continuous chemiluminescence signal which commences upon reaction of the enzyme and rapidly reaches peak intensity. Further modifications and improvements in the light producing reaction can be realized by incorporation of an anionic surfactant and a non-ionic surfactant in combination as will be described in more detail below.



In a preferred method of producing light from the reaction of compound IV with a phosphatase enzyme and a CAC, the reaction is performed at a temperature between 5 °C and 50 °C, usually between 20 °C and 40 °C in an aqueous buffer solution at a pH between 7 and 10.5, preferably between about 8 and 10. It is particularly convenient to conduct the reaction at ambient temperature, without the need to precisely regulate the temperature due to the relatively small influence of temperature on light intensity near ambient temperature. Compound IV is used at a concentration between 1 μM and 20 mM, preferably between 10 μM and 1 mM. The enzyme is preferably an alkaline

phosphatase or an alkaline phosphatase conjugate.

More than one compound of formula I-V can be used concurrently in a method for producing light by the action of a phosphatase enzyme in the presence of a CAC. It can be advantageous in some instances to simultaneously react two or more compounds of formula I-V with the phosphatase enzyme and the CAC. When the two or more compounds have differing luminescent or physical properties, the combination of the two can be desirable to produce a light emitting reaction with characteristics not readily achievable through the use of any one compound. Examples of luminescent and physical properties which can differ between compounds I-V include emission spectrum, duration of light emission, enzyme turnover, rate of rise of emission to maximum, hydrophobicity/hydrophilicity and solubility.

Similarly, more than one CAC can be used concurrently in a method for producing light by the action of a phosphatase enzyme. The CAC is used at a concentration between 10^{-2} M and 10^{-9} M, preferably between about 10^{-4} M and 10^{-7} M. Desirable concentrations of a particular CAC for use in the present methods can be readily determined by means of the methods set forth in the specific examples below

In another aspect, the present invention relates to a reagent composition for producing chemiluminescence by reaction with a phosphatase enzyme comprising an aqueous buffer with a pH between about 7 and about 10.5, a

compound of formula I-V and a CAC in an amount effective to provide increased levels of chemiluminescence.

Additionally, it has been found that incorporation of certain additives into the reaction of a compound of formula I-V and a CAC with a phosphatase enzyme results in further desirable improvements in the light producing reaction. Yet another aspect of the present invention therefore is a reagent composition comprising a compound of formula I-V, a CAC, an effective amount of an anionic surfactant and an effective amount of a non-ionic surfactant. Anionic surfactants serve to substantially increase the speed with which maximum chemiluminescence intensity is reached. Non-ionic surfactants serve to substantially increase the amount or intensity of chemiluminescence which is generated. Use of the latter compositions is particularly advantageous in providing increased light emission which rapidly reaches and maintains peak light intensity.

In contrast to the methods of Applicant's co-pending application U.S. No. 08/585,090 in which cationic surfactants enhance the production of chemiluminescence, the incorporation of cationic surfactants, both monomeric and polymeric, are ineffective in promoting the chemiluminescent reaction which requires the presence of a CAC. In some cases, inclusion of cationic surfactants virtually extinguishes light production. The opposite effect of cationic surfactants in the present methods and compositions underscores the difference in the

chemiluminescent reaction process which results from utilizing a CAC.

Anionic surfactants useful in compositions containing a CAC include alkyl sulfates and alkylsulfonates having an alkyl group of at least ten carbons. A preferred compound is sodium dodecyl sulfate (SDS). Anionic surfactants are preferably used in an amount from about 10 mg/mL to 10 μ g/mL.

Non-ionic surfactants useful in compositions containing a CAC include polyoxyethylenated alkylphenols, polyoxyethylenated alcohols, polyoxyethylenated ethers and polyoxyethylenated sorbitol esters. Preferred non-ionic surfactants include TWEEN 20 and TRITON X-100. Non-ionic surfactants are preferably used in an amount from about 1.0 % to about 0.001 % by weight of the composition.

More extensive lists of exemplary structures of each category of surfactant can be found in any standard treatise on surfactants and are known to those of skill in the art. Numerous representative member surfactants have been tested and found to be effective to varying degrees in increasing the amount or intensity of light produced compared to the amount produced in its absence.

A still further aspect of the present invention is a reagent composition comprising a compound of formula I-V, a CAC, an effective amount of an anionic surfactant, an effective amount of a non-ionic surfactant and an effective amount of a background reducing agent which reduces chemiluminescence produced by the composition in

the absence of a phosphatase enzyme (background chemiluminescence).

Background reducing agents are compounds which reduce chemiluminescence produced by the composition in the absence of a phosphatase enzyme. These agents can also function by preventing the accumulation of background chemiluminescence over a period of time. These agents can also function by improving the ratio of specific signal produced by reaction of the composition with a phosphatase enzyme to background chemiluminescence. Preferred compounds which are effective in reducing the amount of background chemiluminescence include sulfite salts such as lithium sulfite, sodium sulfite and potassium sulfite.

A preferred composition for detection of AP or conjugates in solution comprises an amine buffer, pH 8.5-9, 0.1-1000 μM lucigenin, 0.1-1.0 mM compound 5, 0.1-5 mg/mL SDS, 1-100 $\mu\text{g/mL}$ Na_2SO_3 , 0.01-0.1 % (w/v) TWEEN 20 and 0.1-1.0 mM Mg salt. A preferred composition for detection of AP or conjugates on a membrane comprises an amine buffer, pH 8.5-9, 0.1-1000 μM lucigenin, 0.1-1.0 mM compound 5, 0.1-5 mg/mL SDS, 1-100 $\mu\text{g/mL}$ Na_2SO_3 , 0.01-0.1 % (w/v) TWEEN 20 and 0.1-1.0 mM Mg salt.

Compositions which are effective in the present method can function by increasing the fraction of excited state product molecules which emit light, by increasing the fraction of product molecules which are formed in the excited state, by increasing the rate of reaction or turnover of the enzyme, by increasing the rate of a

subsequent chemical reaction step, by improving the stability of the enzyme, by promoting the association of the enzyme with the compound of formula I, by inhibiting or preventing competitive non-luminescent side reactions
5 or by other as yet unidentified mechanisms.

It is contemplated that fluorescent energy acceptors can be employed to shift the maximum emission to longer wavelengths (red-shifting). Various techniques for red-shifting emission are known in the art of chemiluminescent
10 reactions and assays. The fluorescer can be covalently linked to a compound of formula I-V whereby the excited state of the corresponding compound VI can undergo an intramolecular energy transfer resulting in emission at longer wavelengths. Fluorescers can be added to the
15 reaction solution as separate species. Fluorescers can be linked to an anionic or non-ionic surfactant which can form a micelle in order to bring the fluorescer in close contact to the light emitter. Alternately, the fluorescer can be provided in a non-fluorescent form which is
20 convertible to the fluorescent form by removal of a phosphate group during the enzyme reaction period. Examples of the latter type of compound include fluorescein diphosphate, coumarin phosphates such as 4-methylumbelliferone phosphate and benzothiazole phosphates
25 such as ATTOPHOS (JBL Scientific, San Luis Obispo, California).

An important use of the present chemiluminescent methods is for detecting the presence or amount of an

analyte in an assay procedure by a chemiluminescent reaction. The method comprises the steps of contacting a sample suspected of containing the analyte with a chemiluminescent compound of the present invention and a
5 phosphatase enzyme, detecting the light produced in a qualitative method and, if quantitation is desired, relating the amount of light produced to the amount of the analyte. The relationship between light intensity and amount of analyte can be easily discerned by constructing
10 a calibration curve with known amounts of the analyte. The chemiluminescent compound is typically used in a concentration of about 10^{-5} M to about 10^{-2} M, preferably between about 10^{-4} M and about 10^{-3} M. The phosphatase enzyme is preferably below about 10^{-9} M when detected in a
15 solution. Typical samples which are analyzed by the chemiluminescent reaction method are body fluids such as blood, plasma, serum, urine and semen.

Analyte as used herein means a substance whose presence can be detected or quantified in a sample.

20 Analytes which can be assayed by the present methods include phosphatase enzymes, in which case it would be unnecessary to add additional phosphatase enzyme. The analyte can be an inhibitor of a phosphatase enzyme. The analyte can be any of various classes of organic and
25 biological molecules which can be detected in ligand-binder assays as are generally known in the art and include immunoassays, nucleic acid probe assays, cell receptor assays and the like. In these assays, the analyte

is labeled with a phosphatase enzyme or can be specifically detected through phosphatase-labeled specific binding partners. The phosphatase can be incorporated directly as the label on the analyte binding compound.

- 5 Alternately, the analyte binding compound can be bound to at least one phosphatase-labeled specific binding substance for the analyte binding compound. Alternately, the analyte binding compound can be labeled with at least one second specific binding substance which is then bound
- 10 to a phosphatase-labeled binding partner for the second specific binding substance. The phosphatase enzyme can also be provided as a label on a ligand analog such as an enzyme-hapten conjugate in a competitive assay.

- The phosphatase enzyme which can undergo the
- 15 chemiluminescent reaction include alkaline phosphatase from a bacterial source such as E. coli, or a mammalian alkaline phosphatase or acid phosphatase from plant or mammalian sources. Conjugates of a phosphatase enzyme and a biological molecule can also be used in the method for
- 20 producing chemiluminescence, the only proviso being that the conjugate display phosphatase activity, i.e. the ability to hydrolyze phosphate monoesters. Biological molecules which can be conjugated to one or more molecules of a phosphatase enzyme include DNA, RNA,
- 25 oligonucleotides, antibodies, antibody fragments, antibody-DNA chimeras, antigens, haptens, proteins, lectins, avidin, streptavidin and biotin. Complexes including or incorporating phosphatase enzymes such as

liposomes, micelles, vesicles and polymers which are functionalized for attachment to biological molecules can also be used in the methods of the present invention.

The reaction of a composition of the present invention with a phosphatase enzyme to produce chemiluminescence constitutes a rapid and highly sensitive method for detecting the presence or amount of the phosphatase enzyme. Use of the present method can therefore be made for the purpose of determining the presence or quantity of a phosphatase enzyme in a sample by measuring the amount or intensity of light produced by reaction of the sample with a compound of formula I. Such a determination can be used e.g. in measuring the level of alkaline phosphatase in blood serum as an indication of the status of a patient's liver function or as an index of certain disease conditions. Prostatic acid phosphatase is also useful as a clinical diagnostic index of prostate cancer.

Compositions of the present invention are useful for measurement of acid phosphatase (AcP) produced chemiluminescence which rapidly reaches maximum intensity within 2-5 sec and then decays within a short period of time. Even though the pH of the compositions is far higher than the optimal pH for AcP activity, it is possible to quantitate this enzyme with good sensitivity. Moreover, AcP could be measured in the presence of AP by means of this detection reagent. Since the chemiluminescent signal induced by AcP and AP display different kinetics, it is possible to simultaneously

determine both AcP and AP activities in the same sample in one experiment.

A second area of application for the chemiluminescent measurement of phosphatase activity is in the detection and measurement of enzyme inhibitors. Inhibitors can act reversibly by acting as a substrate in competition with a second substrate such as the compounds of the present invention. Another mode of inhibition known as suicide inhibition acts irreversibly by deactivating the enzyme.

10 Inhibitors of alkaline phosphatase include inorganic phosphate, levamisole and its racemic form tetramisole and other imidazo[1,2-b]thiazoles, L-phenylalanine and L-homoarginine and are identified in R.B. McComb, G.N. Bowers, S. Posen in Alkaline Phosphatase, Plenum Press, New York 1979, pp. 268-275, 332-334, 394-397, 410-413.

15 Inhibitors of acid phosphatase include fluoride, molybdate, orthophosphate ions, tartrate, 4-(fluoromethyl)phenyl phosphate and 4-(fluoromethyl)phenyl phosphonate (J.K. Myers, T.S. Widlanski, Science, 262, 1451-3 (1993)). It is recognized that some substances are only inhibitory at some concentrations and can be only partially inhibitory.

20

Measurement of the quantity or characteristics of an inhibitor, such as the inhibition constant K_i , or half-life for inhibition, $t_{1/2}$, are made by measuring the enzyme activity of a sample containing the enzyme in question in the presence of a substrate producing a detectable product and a quantity of the inhibitor. In a method of detecting

25

a phosphatase inhibitor according to the present invention, a compound of formula I produces light as the detectable product. Reaction of the phosphatase enzyme and chemiluminescent compound is made in the presence and
5 absence of the inhibitor substance and the results are compared to determine the presence or amount of the inhibitor. The effect of the inhibitor can have one or more of any of three effects, a decrease in light intensity, a slower rate of rise of light intensity or a
10 delay period before light emission begins.

A third area of application for the chemiluminescent measurement of phosphatase activity is in gene expression assays. Alkaline phosphatase and in particular an isozyme produced in the placenta which is excreted are useful as
15 reporter genes (J. Alam, J. Cook, Anal. Biochem. 188, 245-54 (1990)). In this type of assay, a gene responsible for expression of a reporter enzyme is cloned into the genetic material of an organism via a plasmid in the vicinity of a promoter or enhancer sequence. The effect of the promoter
20 or enhancer sequence on transcriptional activity is gauged by measuring the level of production of reporter enzyme.

Techniques for performing enzyme assays are well known. With the guidance provided by the examples as taught herein, variations of procedures for preparing
25 samples, determining appropriate quantities and ratios of reagents, reaction times, constructing calibration curves and the like will be within the ability of one of ordinary skill in the art to devise as a matter of routine

experimentation.

Since the reaction is catalyzed by the phosphatase enzyme, exceedingly small quantities of the enzyme are sufficient to produce a detectable amount of light.

5 Sensitivities of 4 zeptomol (4×10^{-21} mol) have been achieved. The ability to detect such small amounts of phosphatase enzymes make the present chemiluminescent technology suitable for analyses of many types of analytes using enzyme-linked assays. Such analyses and assays
10 require the ability to detect small quantities of phosphatase enzymes due to low abundance of the analyte in the sample to be analyzed or to limited sample quantity. In this type of assay, alkaline phosphatase is conjugated to one member of a specific binding pair. An example is a
15 chemiluminescent enzyme-linked immunoassays, such as the so-called enzyme-linked immunosorbent assay or ELISA. Such assays are commonly used in manual format as well as on automated multi-test immunoassay systems. In a typical immunoassay, the analyte hapten, antigen or antibody is
20 assayed by detecting the presence or amount of an enzyme-labeled specific binding partner for the analyte or an enzyme-labeled analog of the analyte. Various assay formats and the protocols for performing the immunochemical steps are well known in the art and do not
25 constitute a part of the invention per se. These assays fall broadly into two categories. Competitive assays feature an immunological binding of a specific antibody with the analyte and an analyte analog, e.g. a detectably

labeled analyte molecule. Sandwich assays result by the sequential or simultaneous binding of two antibodies, one of which is detectably labeled, with the analyte. The detectable enzyme-labeled binding pair so formed can be
5 assayed with the compounds and methods of the present invention. Measurement can be performed with enzyme-labeled species attached to a solid surface or support including beads, tubes, microwells, magnetic particles, test strips, membranes and filters such as are in common
10 use in the art. The detectable enzyme-labeled species can also be present free in solution or enclosed within an organized assembly such as a liposome in which case a lytic agent is employed to lyse the liposome and free the detectable enzyme.

15 Another exemplary use is the detection of proteins by the technique of Western blotting. A sample containing a protein of interest as the analyte is subject to electrophoretic separation. The separated proteins are transferred to a blotting membrane by capillary action or
20 with the aid of an electric field. Such transferred protein is typically detected with a specific primary antibody and an enzyme-labeled secondary antibody which recognizes and binds to the primary antibody. Visualization of marker enzyme activity reflects the
25 presence of the analyte protein. To adapt the methods of the present invention for Western blotting, an AP conjugated secondary antibody can be employed and AP activity measured with chemiluminescence using a compound

of the present invention as the chemiluminescent reagent. Variations on this technique such as using biotinylated antibodies and avidin-AP are considered within the scope of assays able to be performed using the inventive

5 methods.

In addition to the aforementioned antigen-antibody, hapten-antibody or antibody-antibody pairs, specific binding pairs also can include complementary oligonucleotides or polynucleotides, avidin-biotin,
10 streptavidin-biotin, hormone-receptor, lectin-carbohydrate, IgG-protein A, nucleic acid-nucleic acid binding protein and nucleic acid-anti-nucleic acid antibody.


A particularly useful application of the present
15 detection methods is the detection of nucleic acids by the use of enzyme-labeled nucleic acid probes. Methods for analysis and chemiluminescent detection of nucleic acids using enzyme-labels, for example, hybridization assays, DNA detection in Southern blotting, RNA by Northern
20 blotting, DNA sequencing, DNA fingerprinting, colony hybridizations and plaque lifts are all well established techniques. The enzyme label (e.g. AP) can be present as a direct conjugate with a probe oligonucleotide or capture oligonucleotide or it can be incorporated through indirect
25 linking means using art-know methods. Examples of indirect linking means include using hapten-labeled oligonucleotides and anti-hapten-AP conjugates or biotinylated oligonucleotides and avidin-AP conjugates.

Such nucleic acid assays can be performed on a blotting membrane or in solution using oligonucleotides attached to solid surfaces including beads, tubes, microwells, magnetic particles, test strips such as are known in the art.


Process for the Preparation of Compounds I-IV. A

process and intermediates useful for the preparation of chemiluminescent phosphate compounds I-IV of the present invention was disclosed and embodied in the co-pending 585,090 application. For the purposes of elaborating additional exemplary methods and conditions, the following additional explanatory information is presented.

Briefly, the process involves reacting a heterocyclic ester or thioester compound VIII with a base to form an enolate of VIII; phosphorylating the enolate of VIII by reacting the enolate with a phosphorylating agent to form a protected enol phosphate IX and deprotecting the enol phosphate to form the enol phosphate salt compound I by reacting IX with at least one deprotecting agent in the presence of a cationic species M if the cationic species is not a part of the deprotecting agent.




In one embodiment, the protected enol phosphate IX is formed by first reacting the enolate of compound VIII with a phosphorus oxyhalide compound POW_3 , where W is a halogen atom selected from F, Cl, Br and I to form an enol dihalophosphate X. The intermediate enol dihalophosphate X is converted to protected enol phosphate IX by reaction with at least two equivalents of a hydroxylic compound Y-OH. The phosphate triester IX is then converted to phosphate salt I by reaction with at least one deprotecting agent in the presence of a cationic species M if the cationic species is not a part of the deprotecting agent.




20 Preferably the halogen W is Cl. Compounds which can serve as the hydroxylic compound Y-OH include, without limitation, lower alcohols such as methanol and ethanol, substituted lower alcohols such as 3-hydroxypropionitrile ($\text{HOCH}_2\text{CH}_2\text{CN}$) and 2-trimethylsilylethanol, phenol, 25 substituted phenols, benzyl alcohol and others as are generally known.

In another embodiment, the protected enol phosphate IX can be prepared directly from VIII by reacting the enolate

of VIII a phosphorylating agent containing the protecting groups Y and having the formula $W-PO(OY)_2$.



The groups O-Y in this phosphorylating agent can include, :
by way of example, alkoxy such as OCH_3 , OCH_2CH_3 , and the
10 like, substituted alkoxy such as cyanoethoxy ($\text{OCH}_2\text{CH}_2\text{CN}$) or
trimethylsilylethoxy ($\text{OCH}_2\text{CH}_2\text{Si}(\text{CH}_3)_3$), phenoxy,
substituted phenoxy, benzyloxy and others as are generally
known to the skilled organic chemist. The two groups O-Y
can also be combined together as a single group as occurs
15 in the reagent



The enolate is formed by reacting the ester or thioester compound VIII with a strong base in a non-reactive organic solvent at a temperature in the range of -90 to 25 °C. In a preferred mode, the reaction is performed at the temperature of a dry ice/acetone bath, nominally -78 °C for a time and then allowed to warm to 0 to 25 °C for a second time. Suitable solvents are those non-protic solvents which are compatible with the strong base required to form the enolate and include ethers and hydrocarbon solvents. The preferred solvent is

tetrahydrofuran. The strong base and carbonyl compound VIII can be added to the reaction vessel in either order.


The phosphorylation step is performed in the same solution at a temperature in the range of about -78°C to about 25°C . The phosphorylating agent, either POW_3 or $\text{WPO}(\text{OY})_2$, is added in a controlled fashion so as not to cause the reaction solution to become hot. The phosphorylation agent is preferably accompanied by an amine base, preferably pyridine.

10 The deprotection step is accomplished by reacting the protected enol phosphate IX with a deprotecting agent in a quantity sufficient to cause removal of the protecting groups and in the presence of a cationic species M if the cationic species is not a part of the deprotecting agent.

15 Two equivalents of the deprotecting agent are typically required, however for convenience, the deprotecting agent can be used in molar excess. Removal of the protecting groups can in some cases be performed one at a time. For example, in a compound of formula XI wherein the two Y

20 groups together constitute the single group $-\text{CH}_2\text{CH}_2-$, and thereby form a five-membered ring, the first $\text{O}-\text{CH}_2$ bond can be cleaved by treatment with a cyanide salt. The resulting compound XII is further reacted with a base to cleave the second $\text{O}-\text{CH}_2$ bond and generate the salt I.

25





5 The choice of the deprotecting agent will be determined, in part, by the nature of the groups Y to be removed. The deprotecting agent must also not cause undesired side reactions such as hydrolysis of the vinyl
 10 ether or vinyl sulfide group or undesired changes to the heterocyclic group. Preferred deprotecting agents include organic and inorganic bases such as sodium hydroxide, potassium hydroxide, potassium carbonate, sodium
 15 methoxide, sodium ethoxide, potassium t-butoxide, ammonium hydroxide and the like. Other preferred deprotecting agents include nucleophilic agents such as cyanide ion, fluoride ion.

20 To the best of applicant's knowledge, compounds IX, X, XI and XII in which Het is a nitrogen or sulfur-containing heterocyclic group have not been prepared. An enol phosphate ester compound in which an oxygen-containing - lactone ring is appended to the double bond is disclosed in U.S. 3,130,203. This compound however was prepared by a
 25 different process, not involving phosphorylation of an ester or thioester enolate.

EXAMPLES

The preparation and chemiluminescent reaction of the following compounds of formula I are described below for purposes of illustrating the invention in more detail.




5					
10	<u>Compound</u>	<u>R₇ - R₁₄</u>	<u>Z</u>	<u>R₆</u>	
	1	all H	O	phenyl	
	2	all H	O	3,5-difluorophenyl	
	3	R ₉ = OCH ₃	O	phenyl	
	4	R ₉ = Cl	O	2,6-dimethylphenyl	
15	5	all H	S	phenyl	
	6	R ₁₁ -R ₁₂ =	O	phenyl	
					
	7	all H	S	4-fluorophenyl	
20	8	all H	S	4-methoxyphenyl	
	9	all H	S	2,6-dimethylphenyl	
	10	R ₈ , R ₁₃ = F	S	phenyl	
	11	all H	S	trifluoroethyl	
	12	all H	S	4-chlorophenyl	
25	13	all H	S	2-naphthyl	

R₇ - R₁₄ are H unless otherwise indicated.


Each of compounds 1-13 was prepared by the following synthetic scheme. It will be recognized that modifications to the reaction conditions can be made within the confines of the synthetic process described above in order to prepare these and other compounds of formula I.

Scheme 1.



Example 1. Synthesis of Acridan Derivative 1.

5



a. Phenyl acridine-9-carboxylate. Acridine-9-carboxylic acid (1 g, 4.1 mmol) was suspended in thionyl chloride (5 mL) and the reaction mixture was refluxed for 3 h. The solvent was removed under reduced pressure leaving a yellow solid which was dissolved in CH_2Cl_2 and pyridine (350 μL) under argon. This solution was cooled in an ice bath and a solution of phenol (0.78 g, 8.2 mmol) in CH_2Cl_2 was added dropwise. The reaction mixture was stirred overnight at room temperature. After evaporation of solvent, the residue was redissolved in ethyl acetate and washed with water. The organic layer was dried over MgSO_4 and concentrated to obtain a crude material which was chromatographed on silica gel (30 % ethyl acetate/hexane) to yield the pure product as a yellow solid. ^1H NMR (CDCl_3) δ 7.35-7.57 (m, 5H), 7.63-8.37 (m, 8H).

b. Phenyl 10-methylacridinium-9-carboxylate trifluoromethanesulfonate. Phenyl acridine-9-carboxylate (530 mg, 1.7 mmol) was dissolved in CH_2Cl_2 (5 mL) under argon and methyl triflate (1 mL, 8.8 mmol) was added. The solution was stirred overnight at room temperature to yield a thick

yellow precipitate. This precipitate was filtered, washed with ether and dried to obtain the product as yellow crystals. ^1H NMR (acetone- d_6) δ 5.22 (s, 3H), 7.47-7.71 (m, 5H), 8.23-9.07 (m, 8H).

5

c. Phenyl 10-methylacridan-9-carboxylate. Phenyl 10-methylacridinium-9-carboxylate trifluoromethanesulfonate (10 mg, 0.0216 mmol) was suspended in absolute ethanol (10 mL) and the mixture was refluxed for 15 min to obtain a clear solution. Ammonium chloride (88 mg, 1.6 mmol) was added by portions to the solution followed by zinc (108 mg, 1.6 mmol). Addition of zinc caused the yellow color of the solution to disappear immediately. The colorless solution was refluxed for 2 h. TLC of the reaction mixture showed complete conversion to a non polar material. The solution was filtered and precipitate was washed with ethanol (3 x 20 mL). The filtrate was concentrated to obtain an off-white solid which was redissolved in CH_2Cl_2 and washed with water (2 x 15 mL). The organic layer was dried over Na_2SO_4 and concentrated to yield the crude product which was purified by preparative TLC using (30% ethyl acetate:hexane). Pure product was obtained as an off-white solid. ^1H NMR (CDCl_3) δ 3.38 (s, 3H), 5.16 (s, 1H), 6.89-7.37 (m, 13H); ^{13}C NMR (CDCl_3) δ 33.29, 49.72, 112.93, 120.19, 121.36, 125.73, 128.67, 129.16, 129.26, 142.37, 151.04, 170.22.

25

d. 9-(Phenoxyphosphoryloxymethylidene)-10-methyl-

acridan, bis(cyanoethyl) ester. A three-neck flask was
purged with argon and charged with 5 mL of anhydrous THF
and diisopropylamine (0.04 mL, 0.29 mmol). The flask was
cooled in a dry ice-acetone bath. To this solution was
5 added n-butyl lithium (0.116 mL, 0.29 mmol). After 20 min,
a solution of the acridan ester from step (c) (70 mg, 0.22
mmol) in 5 mL of THF was added to this solution and
stirring was continued for 30 min at -78 °C. Finally a
solution of POCl₃ (0.027 mL, 0.29 mmol) and pyridine
10 (0.023 mL, 0.29 mmol) in 3 mL of THF was added and the dry
ice bath was removed. After 45 min, pyridine (0.039 mL,
0.58 mmol) and 3-hydroxypropionitrile (0.094 mL, 1.16
mmol) was added and stirring maintained over night. Then
it was filtered and solvent was removed from the filtrate.
15 The residue was subjected to prep. TLC (80% ethyl acetate/
hexane) to give the pure product; ¹H NMR (CDCl₃) δ 2.35-
2.54 (m, 4H), 3.47 (s, 3H), 3.79-3.90 (m, 2H), 3.98-4.08
(m, 2H), 6.825-7.45 (m, 12H), 7.80-7.83 (dd, 1H); ¹³C NMR
(CDCl₃) δ 19.12, 19.24, 33.63, 62.19, 62.49, 88.72, 92.82,
20 112.40, 112.52, 115.83, 116.07, 119.68, 120.44, 120.71,
123.81, 126.69, 128.03, 128.27, 128.58, 130.09, 142.39,
143.06, 165.73, 202.09.

e. 9-(Phenoxyphosphoryloxymethylidene)-10-methyl-
25 acridan, disodium salt (1). A solution of the
bis(cyanoethyl) phosphate compound (2.897 g, 5.77 mmol) in
50 mL of acetone was purged with Ar for 30 min. An Ar-
purged solution of 479 mg (12 mmol) of NaOH in 7.5 mL of


water was added dropwise and the solution stirred over night. The precipitate which had formed was filtered, washed with 50 mL of Ar-purged acetone and air-dried. The yield was 3.473 g of 1 as a white solid which contained

5 some water. ^1H NMR (D_2O) δ 3.326 (s, 3H), 6.825- 7.45 (m, 11H), 7.80-7.83 (d, 1H); ^{13}C NMR (D_2O) δ 32.95, 102.86, 102.92, 112.30, 115.85, 120.68, 121.01, 122.35, 122.41, 122.62, 127.48, 127.66, 128.23, 129.66, 143.17, 143.32, 144.66, 156.01; ^{31}P NMR (D_2O) δ 0.581 (rel. to ext. H_3PO_4).

10

Example 2. Synthesis of Acridan Derivative 2.

15



a. 3',5'-Difluorophenyl acridine-9-carboxylate.

20 Acridine-9-carboxylic acid (0.25 g) was suspended in 10 mL thionyl chloride and the reaction mixture was refluxed for 3 h. The solvent was removed under reduced pressure to obtain a yellow solid which was dissolved in CH_2Cl_2 and pyridine (0.22 g) under argon. A solution of 3,5-difluoro-

25 phenol (0.16 g) in CH_2Cl_2 was added dropwise. The solution was stirred overnight at room temperature then diluted with more CH_2Cl_2 (100 mL) and washed with water (3 x 50 mL). The organic layer was dried over Na_2SO_4 and

concentrated to obtain the product which was further purified by chromatography on silica gel (30 % ethyl acetate/hexane) to yield the pure product as a creamy solid. ^1H NMR (CDCl_3) δ 6.84-7.09 (m, 3H), 7.67-8.37 (m, 8H).

b. 3',5'-Difluorophenyl 10-methylacridinium-9-carboxylate trifluoromethanesulfonate. The acridine ester from step (a) (0.20 g) was dissolved in CH_2Cl_2 (5 mL) under argon and methyl triflate (0.472 mL 7 eq.) was added. The solution was stirred overnight at room temperature to yield a thick yellow precipitate. This precipitate was filtered, washed with ether and dried to obtain the product as yellow crystals. ^1H NMR (acetone-d_6) δ 5.25 (s, 3H), 7.22-7.59 (m, 3H), 8.23-9.09 (m, 8H).


c. 3',5'-Difluorophenyl 10-methylacridan-9-carboxylate. The acridinium ester from step (b) (0.10 g) was dissolved in 10 mL of glacial acetic acid to obtain a yellow solution and zinc was added (1.30 g) causing immediate decolorization of the solution. After 5 min - stirring at room temperature, TLC of the reaction mixture showed a nonpolar material. The acetic acid was decanted and the solid washed with CH_2Cl_2 . The combined organic solutions were evaporated to obtain a crude solid which was redissolved in CH_2Cl_2 and washed with 2 or 3 - 50 mL portions of water. The crude material obtained after evaporation of CH_2Cl_2 was chromatographed on silica gel

(20-30 % ethyl acetate/ hexane) to yield the pure product as a white solid. ^1H NMR (CDCl_3) δ 3.44 (s, 3H), 5.16 (s, 1H), 6.49-6.65 (m, 3 H), 6.99-7.36 (m, 8H).

- 5 d. 9-(3,5-Dimethylphenoxy)phosphoryloxymethylidene)-
10-methylacridan, bis(cyanoethyl) ester. A three-neck flask was purged with argon and charged with 4 mL of anhydrous THF and diisopropylamine (0.0494 mL, 0.35 mmol). The flask was cooled in a dry ice-acetone bath. To this
10 solution was added n-butyllithium (0.141 mL, 0.35 mmol). After 20 min, a solution of the acridan ester from step (c) (75 mg, 0.22 mmol) in 4 mL of THF was added dropwise to the LDA. The funnel was washed with an additional 4 mL of Ar-purged THF which was added to the reaction solution.
15 Stirring was continued for 30 min at -78°C . Finally a solution of POCl_3 (0.034 mL, 0.35 mmol) and pyridine (0.057 mL, 0.35 mmol) in 4 mL of THF was added causing the brown solution to decolorize. The dry ice bath was removed and stirring continued for 15 min. A solution of pyridine
20 (0.095 mL, 1.17 mmol) and 3-hydroxypropionitrile (0.080 mL, 1.17 mmol) in 4 mL of THF was added and stirring maintained over night. Then it was filtered and the solvent was removed from the filtrate. The residue was subjected to prep. TLC (55 % ethyl acetate/hexane) from
25 which a small amount of the pure product was isolated; ^1H NMR (acetone- d_6) δ 2.82 (m, 4H), 3.51 (s, 3H), 4.10-4.31 (m, 4H), 6.75-7.93 (m, 11H).

e. 9-(3,5-Dimethylphenoxy)phosphoryloxymethylidene)-10-methylacridan, disodium salt (2). A solution of the bis(cyanoethyl)phosphate from step (d) 5 mg, 5.77 mmol) in 5 mL of methanol was purged with argon for 30 min with ice cooling. An Ar-purged solution of 6 mg of Na_2CO_3 in 0.5 mL of water was added dropwise and the solution stirred under argon over a weekend. The solvent was evaporated and the solid compound 2 dried under vacuum. ^1H NMR (D_2O) δ 3.34 (s, 3H), 6.66-8.09 (m, 11H).

Example 3. Synthesis of Acridan Derivative 3.



3

(E/Z mixture)

a. N-Methyl-3-methoxyisatin. 3-Methoxydiphenylamine (250.2 g, 1.26 mol) dissolved in 900 mL of CH_2Cl_2 was added to a hot solution of 140 mL (1.6 mol) of oxalyl chloride in 1 L of CH_2Cl_2 under an argon atmosphere. The addition was completed over a 2 hour period. Heating was discontinued during the addition since the reaction produced enough heat to maintain reflux. Stirring was continued at reflux temperature for another 30 min after adding the amine. The resulting brown solution was

evaporated to remove excess oxalyl chloride. The brown solid was redissolved in 1.8 L of CH_2Cl_2 , which was purged with argon. Aluminum chloride (359 g, 2.69 mol) was added in portions to the solution over a 45 min period while the mixture was stirred mechanically. The mixture became thick during the addition and began to reflux. Reflux was maintained for an additional hour after all of the AlCl_3 had been added. The cooled mixture was evaporated in vacuo and quenched with 4 kg of ice, ca. 500 mL of water and 240 mL of conc. HCl. The dark solid was carefully crushed to small particles while stirring at ice temperature for 2 h. The solid was then filtered off with suction and washed with a total of 8 L of water. Air drying produced an orange solid which was a mixture of isomeric isatins containing > 90% of one isomer.

b. 3-Methoxyacridine-9-carboxylic acid. The isatin mixture from step (a) was refluxed in a 10% KOH solution in water for ca. 40 h. The solution was cooled to < 60 °C and filtered through glass wool into a mixture of 480 mL of conc. HCl and 10 L of ice water which was stirred vigorously. The solid was suction filtered after 1 h and then washed with 8 L of water. The solid was air-dried over night and then dried under vacuum. ^1H NMR ($\text{CD}_3\text{OD}/\text{NaOH}$) δ 4.062 (s, 3H), 7.25-8.18 (m, 7H).

c. Phenyl 3-methoxyacridine-9-carboxylate. 3-Methoxyacridine-9-carboxylic acid (0.50 g) was suspended in 10 mL

thionyl chloride (3-10 mL) and reaction mixture was refluxed for 3 h. The solvent was removed under reduced pressure to obtain a yellow solid which was dissolved in CH_2Cl_2 and pyridine (0.44 g) under argon. A solution of
5 2,6-difluorophenol (0.32 g) in CH_2Cl_2 was added dropwise. The solution was stirred overnight at room temperature then diluted with more CH_2Cl_2 (100 mL) and washed with water (3 x 50 mL). The organic layer was dried over Na_2SO_4 and concentrated to obtain the product which was further
10 purified by chromatography on silica gel (30 % ethyl acetate/hexane) to yield the pure product as a creamy solid. ^1H NMR (acetone- d_6) δ 4.08 (s, 3H), 7.39-8.29 (m, 12H).

15 d. Phenyl 3-methoxyacridan-9-carboxylate. Ammonium chloride (3.45 g, 64 mmol) and 4.2 g of zinc (64 mmol) were added to an Ar-purged suspension of 0.85 g (2.5 mmol) of the ester from step (c) in 50 mL of ethanol. The mixture was stirred for 2 h, filtered and the solids
20 washed with CH_2Cl_2 . The combined solutions were evaporated to dryness producing a white solid which was dissolved in ethyl acetate and washed with 3 x 25 mL of water. The product was used without further purification. ^1H NMR (CDCl $_3$) δ 3.76 (s, 3H), 5.29 (s, 1H), 6.47-7.36 (m, 12H),
25 8.21 (br s, 1H).

e. Phenyl 3-methoxy-10-methylacridan-9-carboxylate.
The acridan compound from the previous step (0.77 g, 2.3

mmol) was dissolved in CH_2Cl_2 and methyl triflate (3.8 g, 23 mmol) was added. TLC indicated complete methylation after stirring over night. The volatiles were evaporated and the residue purified by chromatography using 18% ethyl acetate / hexane to elute the column. The product contained .75 g of a sticky solid. ^1H NMR (CDCl_3) δ 3.41 (s, 3H), 3.84 (s, 3H), 5.13 (s, 1H), 6.52-7.38 (m, 12H).

f. (E,Z)-9-(Phenoxy)phosphoryloxymethylidene)-3-methoxy-10-methylacridan, bis(cyanoethyl) ester. A solution of LDA was prepared essentially as described in Example 2d. After stirring 25 min at -78°C , a solution of the acridan ester from step 3e (95 mg, 0.26 mmol) in 4 mL of THF was added dropwise. Stirring was continued for 30 min at -78°C . Finally a solution of POCl_3 (0.04 mL) and pyridine (0.05 mL) in 4 mL of THF was added causing the yellow solution to decolorize. The dry ice bath was removed and stirring continued for 105 min. The solution was cooled in an ice bath and treated with pyridine (0.106 mL) and 3-hydroxypropionitrile (0.090 mL) in 4 mL of THF. After stirring for 20 h at room temperature, the reaction mixture was filtered and the yellow precipitate washed with THF. The reaction solvent was evaporated in vacuo and the residue combined with the precipitate. This material was subjected to prep. TLC (60 % ethyl acetate/hexane) from which a small amount of the product was isolated as a mixture of double bond isomers; ^1H NMR (acetone- d_6) δ 2.75-2.82 (m, 4H), 3.52 (s, 3H), 3.82 (s), 3.91 (s), 4.00-4.26


(m, 4H), 6.47-7.96 (m, 12H).

g. (E,Z)-9-(Phenoxy)phosphoryloxymethylidene)-3-methoxy-10-methylacridan, disodium salt (3). The

5 bis(cyano-ethyl)phosphate from step 3(f) was dissolved in 10 mL of MeOH purged with Ar and cooled in an ice bath. An Ar-purged solution of 11 mg of Na_2CO_3 in 1 mL of water was added dropwise and the solution stirred for 19 h. Another 3 mL of methanol was added and stirring continued for 10 another 24 h to allow the deprotection to go to completion. The solvent was evaporated, the solid compound 3 washed with CH_2Cl_2 and dried under vacuum. ^1H NMR (D_2O) δ 3.24 (s, 3H), 3.66 (s), 3.83 (s), 6.35-8.07 (m, 12H).

15 Example 4. Synthesis of Acridan Derivative 4.

20



(E/Z mixture)

a. 3-Chloroisatin. 3-Chlorodiphenylamine (20 g, 0.98 mol) dissolved in 250 mL of CH_2Cl_2 was added to a solution of 14.0 g (0.11 mol) of oxalyl chloride in 150 mL of CH_2Cl_2 under an argon atmosphere. The solution was refluxed for 30 min. The resulting solution was evaporated to remove

excess oxalyl chloride. The solid was redissolved in CH_2Cl_2 which was purged with argon. Aluminum chloride (54.48 g) was added in portions to the solution over a 45 min period while the mixture was stirred mechanically. The mixture
5 became thick during the addition and began to reflux. Reflux was maintained for an additional hour after all of the AlCl_3 had been added. The cooled mixture was evaporated in vacuo and quenched with 900 mL of 1M HCl in an ice bath. The solid was then filtered off with suction
10 and washed with water. ^1H NMR (CDCl_3) δ 6.8-7.8 (m, 8H).

b. 3-Chloroacridinecarboxylic acid. The isatin product from step (a) was refluxed in 300 mL of 10% KOH solution in water for ca. 48 h. The solution was cooled
15 and acidified to pH 2-3 with HCl. The yellow solid was suction filtered after 1 h and then washed with water. The solid was air-dried over night, washed with CH_2Cl_2 and then air-dried yielding 23.76 g of the acridine acid. ^1H NMR ($\text{DMSO}-d_6$) δ 7.72-8.30 (m, 7H).

20

c. 2',6'-Dimethylphenyl 3-chloroacridinecarboxylate. The acid (3.0 g) was esterified with 2,6-dimethylphenol (2.24 g, 1.5 eq.) and pyridine (1.98 mL, 2 eq.) in ca. 100 mL of CH_2Cl_2 at room temperature under nitrogen. The
25 solvent was evaporated and the residue purified by column chromatography using 10-30% ethyl acetate/hexane which allowed the separation of the 1-chloro- (minor) and 3-chloro- isomers; ^1H NMR (CDCl_3) δ 2.430 (s, 6H), 7.20-7.28

(m, 3H), 7.60-7.75 (m, 3H), 8.28-8.50 (m, 4H).

d. 2',6'-Dimethylphenyl 3-chloro-10-methylacridinium-carboxylate trifluoromethanesulfonate. The acridine

5 compound from the previous step (2.47 g, 6.7 mmol) was dissolved in CH₂Cl₂ and methyl triflate (3.8 mL, 5 eq.) was added. TLC indicated complete methylation after stirring over night. The volatiles were evaporated and the residue purified by chromatography using 5% ethyl acetate/hexane
10 to elute the column. The product was recrystallized from CH₂Cl₂/hexane yielding 1.7 g of white crystals.

e. 2',6'-Dimethylphenyl 3-chloro-10-methylacridan-carboxylate. Ammonium chloride (2.91 g, 64 mmol) and 3.56
15 g of zinc (64 mmol) were added to an Ar-purged suspension of 2.00 g (5.4 mmol) of the ester from step (c) in 50 mL of ethanol. The mixture was stirred for 2 h, filtered and the solids washed with CH₂Cl₂. The combined solutions were evaporated to dryness producing a white solid. The product
20 was used without further purification. ¹H NMR (CDCl₃) δ 1.746 (s, 6H), 3.400 (s, 3H), 5.152 (s, 1H), 6.88-7.04 (m, 7H), 7.26-7.39 (m, 3H).

f. (E,Z)-9-(2,6-Dimethylphenoxy)phosphoryloxymethylidene)-3-chloro-10-methylacridan, bis(cyanoethyl) ester.
25

The acridan ester (0.100 g, 0.29 mmol) from step (e) was added to a solution of LDA (0.44 mmol) in 10 mL of THF. After stirring 30 min at -78 C, a solution of POCl₃ (42

μL) and pyridine (47 μL) in 4 mL of THF was added. The dry ice bath was removed and stirring continued for 30 min.

TLC (30% ethyl acetate/hexane) showed that the starting material was completely reacted. The solution was cooled

5 in an ice bath and treated with pyridine (117 μL) and 3-hydroxypropionitrile (94 μL) in 4 mL of THF. After stirring over night at room temperature, the precipitated was filtered away and the reaction solvent evaporated. The residue was subjected to prep. TLC (60 % ethyl

10 acetate/hexane) from which the isomeric products was isolated; ¹H NMR (CD₃OD) δ (1.91 (s), 1.95 (s), combined 3H), (2.03 (s), 2.04 (s), combined 3H), 2.69-2.76 (m, 4H), (3.52 (s), 3.59 (s), combined 3H), 4.16-4.34 (m, 4H), 6.97-7.65 (m, 10H).


15

g. (E,Z)-9-(2,6-Dimethylphenoxy)phosphoryloxymethylidene)-3-chloro-10-methylacridan, disodium salt (4). The bis(cyanpethyl)phosphate from the step above is dissolved in 10 mL of methanol purged with argon and then cooled in an ice bath. An Ar-purged solution of 11 mg of Na₂CO₃ in 1 mL of water is added dropwise and the solution stirred under argon until the deprotection has gone to completion. The solvent is evaporated, the solid washed with CH₂Cl₂ and dried under vacuum to produce 4.

25

Example 5. Synthesis of Acridan Derivative 5.

5



5

a. 9-(Phenythiophosphoryloxymethylidene)-10-methyl-acridan, bis(cyanoethyl) ester. Phenyl 10-methylacridan-

10 9-thiocarboxylate (70 mg, 0.2 mmol) was added to a solution of LDA (0.24 mmol) in THF. After stirring 30 min at -78 C, a solution of POCl₃ (25 μL) and pyridine (21 μL) in 4 mL of THF was added. The dry ice bath was removed and stirring continued for 30 min. TLC (30% ethyl

15 acetate/hexane) showed that the starting material was completely reacted. The solution was cooled in an ice bath and treated with pyridine (210 μL) and

3-hydroxypropionitrile (44 μL) in 4 mL of THF. After stirring over night at room temperature, the precipitated

20 pyridine-HCl was filtered away and the reaction solvent evaporated in vacuo. The residue was subjected to prep. TLC (ethyl acetate) from which the product was isolated;


¹H NMR (CDCl₃) δ 2.4-2.6 (m, 4H), 3.521 (s, 3H), 3.8-4 (m, 2H), 4.0-4.1 (m, 2H), 6.9-7.2 (m, 4H), 7.22-7.5 (m, 7H),

25 7.75-8 (dd, 2H).

b. 9-(Phenylthiophosphoryloxymethylidene)-10-methyl-acridan, disodium salt (5). A solution of the

bis(cyanoethyl) phosphate compound (0.59 g, 1.21 mmol) in 50 mL of acetone was purged with argon for 30 min. An Ar-purged solution of 104 mg (2.6 mmol) of NaOH in 10 mL of water was added dropwise and the solution stirred under argon over night. The precipitate which had formed was suction filtered, washed with 50 mL of Ar-purged acetone and air-dried. The yield was 0.50 g of **5** as a slightly yellow solid which contained a slight amount of acetone. ¹H NMR (D₂O) δ 3.36 (s, 3H), 6.9-7.4 (m, 11H), 7.75-7.8 (d, 1H), 8.2-8.22 (d, 1H) ; ³¹P NMR (D₂O) δ 1.85 (rel. to ext. H₃PO₄).

Example 6. Synthesis of Benz[clacridan Derivative 6.



6

(E/Z mixture)

a. Benz[clacridine-7-carboxylic acid. 1-Naphthyl-phenylamine (10 g, 46 mmol) dissolved in 25 mL of CS₂ was refluxed with 4.8 mL (55 mmol) of oxalyl chloride under argon. The resulting brown solution was evaporated to remove excess oxalyl chloride. The brown solid was dissolved in 60 mL of CS₂ which was purged with argon. AlCl₃ (21.47 g, 160 mmol) was added in portions to the solution while the mixture was stirred. The mixture became thick during the addition and began to reflux. Reflux was

maintained for two more hours after all of the AlCl_3 had been added. The cooled mixture was evaporated in vacuo and quenched with 1 M HCl and ice. The dark solid was carefully crushed to small particles and the solid
5 filtered off with suction and washed with water.

The isatin product was refluxed in a 10% KOH solution in water over night. The solution was cooled to $< 60^\circ\text{C}$ and neutralized with 6 M HCl and ice. The solid was suction filtered, washed with water and air-dried. ^1H NMR
10 ($\text{DMSO}-d_6$) δ 7.80-8.13 (m, 8H), 8.36-8.39 (d, 1H), 9.38-9.40 (m, 1H).

b. Phenyl benz[clacridine-7-carboxylate.

Benz[c]acridine-7-carboxylic acid (6.0 g) was suspended in
15 75 mL thionyl chloride (3-10 mL) and reaction mixture was refluxed for 4 h. The solvent was removed under reduced pressure and the product dissolved in 75 mL of CH_2Cl_2 . Pyridine (8.9 mL) and phenol (2.27 g) were added and the solution was stirred for 3 days at room temperature under
20 Ar. Purification was effected by chromatography on silica gel (50 % CH_2Cl_2 /hexane) to yield 3.05 g of the pure product as a light brown solid. ^1H NMR (CDCl_3) δ 7.40 (t, 1H), 7.48-7.59 (m, 4H), 7.71-8.01 (m, 7H), 8.259 (d, 1H, $J=7.8$ Hz), 8.465 (d, 1H, $J=7.8$ Hz), 9.54-9.57 (m, 1H).

25

c. Phenyl benz[clacridan-7-carboxylate. Ammonium chloride (10 g) and 10 g of zinc were added to an Ar-purged suspension of 1.0 g of the ester from step (b) in

ethanol. The mixture was stirred for 2 h, filtered and the solids washed with CH_2Cl_2 . The combined solutions were evaporated to dryness producing a white solid which was dissolved in ethyl acetate and washed with 3 x 25 mL of water. The product was purified by chromatography using 10 % ethyl acetate/hexane. ^1H NMR (CDCl_3) δ 5.48 (s, 1H), 5.69-7.27 (m, 9H), 7.41-7.55 (m, 5H), 7.83-7.88 (m, 2H).

d. Phenyl 12-methylbenz[clacridan-7-carboxylate]. The acridan compound from the previous step (0.70 g) was dissolved in CH_2Cl_2 and methyl triflate (2.3 mL) was added. TLC indicated complete methylation after stirring over night. The volatiles were evaporated and the residue purified by chromatography using 5 % ethyl acetate/hexane to elute the column. ^1H NMR (CDCl_3) δ 3.75 (s, 3H), 5.21 (s, 1H), 6.92-6.94 (d, 2H), 7.09-7.62 (m, 11H), 7.86-7.89 (dd, 1H), 8.23-8.26 (d, 1H); ^{13}C NMR (CDCl_3) δ 45.07, 49.99, 120.01, 121.23, 122.66, 123.51, 124.69, 125.30, 125.45, 125.84, 126.33, 126.97, 128.39, 128.64, 129.36, 134.89, 141.20, 147.09, 151.00, 170.89.

e. (E,Z)-9-(Phenoxy)phosphoryloxymethylidene)-12-methylbenz[clacridan, bis(cyanoethyl) ester]. A solution of LDA was prepared as described above. After stirring 25 min at -78 C, a solution of the benzacridan ester (0.30 g) in THF was added dropwise. Stirring was continued for 30 min at -78 °C. Finally a solution of POCl_3 (0.123 mL) and pyridine (0.107 mL) in THF was added causing the deep red


solution to become orange. The dry ice bath was removed and stirring continued for 1 h. The solution was cooled in an ice bath and treated with pyridine (0.213 mL) and 3-hydroxypropionitrile (0.180 mL) in THF. After stirring
5 over night at room temperature, the reaction mixture was filtered and the precipitate washed with THF. The reaction solvent was evaporated in vacuo and the residue combined with the precipitate. This material was subjected to prep. TLC (60 % ethyl acetate/hexane) from which a small amount
10 of the product was isolated as a mixture of double bond isomers; ^1H NMR (CDCl_3) δ 2.44-2.52 (m, 4H), 3.81-4.10 (m, 7H), 6.89-8.24 (m, 15H).

f. (E,Z)-9-(Phenoxy)phosphoryloxymethylidene)-12-
15 methylbenz[clacridan, disodium salt (6)]. The bis(cyano-ethyl)phosphate from the step above was dissolved in 10 mL of acetone purged with argon and then cooled in an ice bath. An Ar-purged solution (0.1 mL of 2M NaOH in water) was added dropwise and the solution stirred under Ar over
20 night. The solid was filtered and the product compound 6 was washed with acetone and dried. ^1H NMR (D_2O) δ 3.52 (s, 3H), 6.85-6.90 (t, 1H), 7.03-7.06 (d, 2H), 7.15-7.23 (m, 3H), 7.33-7.48 (m, 5H), 7.72-7.75 (d, 1H), 7.87-7.89 (d, 1H), 8.10-8.17 (t, 2H).

25

Example 7. Synthesis of Acridan Derivative 7.

5



7

a. 4'-Fluorophenyl acridine-9-thiocarboxylate.

- 10 Acridine-9-carboxylic acid (10 g, 45 mmol) was suspended in thionyl chloride (ca. 100 mL) and the reaction mixture was refluxed for 2 h. The solvent was removed under reduced pressure leaving a yellow solid which was dissolved in 100 mL of CH₂Cl₂ under argon. 4-
- 15 Fluorothiophenol (5.25 mL, 49 mmol) was added followed by 30 mL of pyridine. The reaction mixture became reddish brown and warmed nearly to reflux. The mixture was stirred for 2 days at room temperature under argon. After evaporation of solvent, the residue was washed with 800 mL
- 20 of hexane, and then with a total of 1 L of water. The remaining solid was dissolved in 300 mL CH₂Cl₂ and passed through a plug of Na₂SO₄/ silica gel/Na₂SO₄. The thioester was obtained (11.55 g) as a yellow solid. ¹H NMR (CDCl₃) δ 7.20 (t, 2H), 7.60-7.68 (m, 4H), 7.80-7.88 (m, 2H), 8.129
- 25 (d, 2H), 8.278 (d, 2H).

b. 4'-Fluorophenyl 10-methylacridinium-9-thiocarboxylate trifluoromethanesulfonate. 4'-Fluorophenyl acridine-

9-thiocarboxylate (11.36 g, 34 mmol) was dissolved in CH_2Cl_2 (100 mL) under argon and methyl triflate (20 mL) was added. The solution became brown and produced a yellow precipitate. After 2 days, the precipitate was filtered, washed with 50 mL of CH_2Cl_2 and 500 mL of hexane and dried to obtain the product as yellow solid (15.3 g). ^1H NMR ($\text{DMSO}-d_6$) δ 4.93 (s, 3H), 7.53 (t, 2H), 7.97-8.37 (m, 2H), 8.14 (t, 2H), 8.54 (t, 2H), 8.61 (d, 2H), 8.92 (d, 2H).

10 c. 4'-Fluorophenyl 10-methylacridan-9-thiocarboxylate.

With room lighting off, 4'-fluorophenyl 10-methyl-acridinium-9-thiocarboxylate trifluoromethanesulfonate (15.05 g, 30 mmol) was suspended in 2-propanol (150 mL) and 2.1 mL of acetic acid. The solution was purged with argon for 30 min. and then 10.0 g of zinc powder added. Addition of zinc caused the yellow color of the solution to disappear within minutes. The mixture was stirred over night. The mixture was filtered and the solids washed with 2-propanol and then hexane. The remaining solid was washed with CH_2Cl_2 . Since a TLC analysis showed that the product had dissolved in both solutions, the filtrates were concentrated and combined to obtain a brown solid which was redissolved in CH_2Cl_2 and passed through a plug of Na_2SO_4 /silica gel/ Na_2SO_4 , eluting with CH_2Cl_2 . A mixture of the expected acridan thioester and an acridan isopropyl ester resulting from transesterification was obtained. The two products were separated by column chromatography (50% CH_2Cl_2 :hexane). One fraction of pure acridan thioester


product (2.5 g) was obtained as a white solid. A second fraction was obtained as a mixture with the isopropyl ester. ^1H NMR (CDCl_3) δ 3.45 (s, 3H), 5.08 (s, 1H), 6.94-7.06 (m, 6H), 7.17-7.24 (m, 2H), 7.31-7.39 (m, 4H).

5

d. 9-(4-Fluorophenylthiophosphoryloxymethylidene)-10-methylacridan, bis(cyanoethyl) ester. With room lighting off, 4-fluorophenyl 10-methylacridan-9-thiocarboxylate (1.00 g, 2.87 mmol) in 10 mL of anhydrous THF was added dropwise to a solution of LDA in THF at -78°C . After stirring 60 min at -78°C , the yellow solution was treated with a solution of POCl_3 (0.79 g) and pyridine (3.0 mL) in 10 mL of THF over a 25 minute period. The dry ice bath was removed after an additional 15 hour and stirring continued for 2 h. The solution was cooled in an ice bath and treated with 3-hydroxypropionitrile (0.71 mL) and the ice bath removed after the addition was complete. After stirring over night at room temperature, the precipitated pyridine-HCl was 20 filtered off and washed with 50 mL of THF. The combined filtrates were evaporated in vacuo. The residue was separated by column chromatography using 50-100 % ethyl acetate/hexanes from which the 0.658 g of the product was isolated; ^1H NMR (CDCl_3) δ 2.44-2.64 (m, 4H), 3.49 (s, 3H), 25 3.86-4.16 (m, 4H), 6.92-7.15 (m, 6H), 7.26-7.46 (m, 4H), 7.786 (d, 1H), 7.906 (d, 1H); ^{31}P NMR (CDCl_3) δ -9.49 (p) (rel. to ext. H_3PO_4).

e. 9-(4-Fluorophenylthiophosphoryloxymethylidene)-10-methylacridan, disodium salt (7). With room lighting off, a solution of the bis(cyanoethyl) phosphate compound (0.658 g) in 20 mL of acetone under argon for 30 min. A solution of 116.5 mg of NaOH in 4.0 mL of water was added and the solution stirred under argon over night. The precipitate which had formed was suction filtered, washed with 100 mL of acetone and air-dried. A second crop of solid formed in the filtrate and was combined with the first crop. The yield was 0.545 g of **7** as a slightly yellow solid. ^1H NMR (D_2O) δ 3.16 (s, 3H), 6.80-6.98 (m, 4H), 7.06-7.14 (m, 4H), 7.21 (t, 1H), 7.32 (t, 1H), 7.78 (d, 1H), 8.19 (d, 1H); ^{31}P NMR (D_2O) δ 1.22 (s) (rel. to ext. H_3PO_4).

Example 8. Synthesis of Acridan Derivative 8.



8

a. 4'-Methoxyphenyl acridine-9-thiocarboxylate. The Acridine-9-carbonyl chloride (2.0 g), made as described above was dissolved in 20 mL of CH_2Cl_2 . 4-Methoxythiophenol (1.27 g) was added dropwise followed by 1.96 g of pyridine. The reaction mixture was stirred for 3.5 days at

room temperature. The precipitate was collected, washed with water and dried to produce a first crop of product. The CH_2Cl_2 solution was washed with 3 x 50 mL of water, dried and evaporated to obtain a crude material which was chromatographed on silica gel (ethyl acetate) to yield a second crop of the pure product. ^1H NMR (CDCl_3) δ 3.86 (s, 3H), 7.01 (d, 1H), 7.56 (d, 2H), 7.62-8.28 (m, 8H).

b. 4'-Methoxyphenyl acridan-9-thiocarboxylate. 4'-Methoxyphenyl acridine-9-thiocarboxylate (2.0 g) was suspended in 2-propanol (150 mL). NH_4Cl (7.35 g) was added by portions to the solution followed by zinc (8.9 g). The solution was stirred at room temperature for 2 h and then warmed gently for 3 h. TLC of the reaction mixture showed complete consumption of the starting material. The solution was filtered and precipitate was washed with CH_2Cl_2 . The filtrate and washings were concentrated to obtain a crude product which was redissolved in CH_2Cl_2 and washed with water (3 x 50 mL). The organic layer was dried and concentrated to yield the product which contained a small amount of the acridan isopropyl ester. ^1H NMR (CDCl_3) δ 3.74 (s, 3H), 5.19 (s, 1H), 6.29 (br s, 1H), 6.75-7.31 (m, 12H).

c. 4'-Methoxyphenyl 10-methylacridan-9-thiocarboxylate trifluoromethanesulfonate. 4'-Methoxyphenyl acridan-9-thiocarboxylate (1.8 g) was dissolved in CH_2Cl_2 under argon and methyl triflate (6.3 g) was added. The solution was


stirred overnight at room temperature to effect complete methylation. The reaction mixture was concentrated and the residue separated by column chromatography (30 % ethyl acetate/hexane) to obtain the 1.08 g of the product. ¹H NMR (CDCl₃) δ 3.45 (s, 3H), 3.76 (s, 3H), 5.07 (br s, 1H), 6.81-7.36 (m, 12H).

d. 9-(4-Methoxyphenylthiophosphoryloxymethylidene)-10-methylacridan, bis(cyanoethyl) ester. A solution of 4-methoxyphenyl 10-methylacridan-9-thiocarboxylate (700 mg) in 10 mL of dry THF was added dropwise to a solution of LDA (1.4 eq.) in 10 mL of dry THF maintained at -78 C. After stirring for 1 h at -78 C, a solution of POCl₃ (520 mg) and pyridine (1.64 g) in 4 mL of THF was added dropwise. Stirring continued for 30 min at -78 °C, the dry ice bath was removed and stirring continued for another hour. The solution was again cooled in an ice bath and pyridine (1 mL) and 3-hydroxypropionitrile (0.86 mL) added dropwise. After stirring over night at room temperature, the precipitated pyridine-HCl was filtered away and the reaction solvent evaporated in vacuo. The residue was taken up in ethyl acetate and washed with water (3 x 25 mL). The organic solution was dried and concentrated, the crude product subjected to column chromatography (30-100 % ethyl acetate/hexane) from which the product was isolated as a light yellow solid; ¹H NMR (CDCl₃) δ 2.44-2.62 (m, 4H), 3.51 (s, 3H), 3.82 (s, 3H), 3.88-4.11 (m, 4H), 6.53-7.41 (m, 10H), 7.84-7.91 (m, 2H); ³¹P NMR (CDCl₃) δ -

9.63 (p) (rel. to ext. H_3PO_4).

e. 9-(4-Methoxyphenylthiophosphoryloxymethylidene)-10-methylacridan, disodium salt (8). A solution of the bis(cyanoethyl) phosphate compound (0.42 g) in 11 mL of acetone was purged with argon. A 614 μL portion of a 2.5 M NaOH solution in water was added dropwise and the solution stirred under argon over night. The precipitate which had formed was suction filtered and air-dried producing a quantitative yield of **8** as a slightly yellow solid. ^1H NMR (D_2O) δ 3.32 (s, 3H), 3.70 (s, 3H), 6.69-7.33 (m, 10H), 7.80 (d, 1H), 8.18 (d, 1H).

Example 9. Synthesis of Acridan Derivative 9.



9

10 a. 2',6'-Dimethylphenyl acridine-9-thiocarboxylate.

Acridine-9-carboxylic acid (2 g) was suspended in thionyl chloride (25 mL) and the reaction mixture was refluxed for 2.5 h. The solvent was removed under reduced pressure leaving a solid which was dissolved in 50 mL of CH_2Cl_2 .

15 This solution was cooled in an ice bath and 2,6-dimethyl-

thiophenol (1.23 g) was added dropwise followed by 3.6 mL of pyridine. The reddish brown reaction mixture was stirred for 1.5 h at room temperature. The reaction mixture was diluted with CH_2Cl_2 and washed with 3 x 50 mL of water. The organic layer was dried and concentrated to obtain an orange product which was chromatographed on silica gel (ethyl acetate) to yield 2.5 g of the pure product as an off-white solid. ^1H NMR (CDCl_3) δ 2.68 (s, 6H), 7.30-7.41 (m, 3H), 7.63-8.31 (m, 8H).

b. 2',6'-Dimethylphenyl acridan-9-thiocarboxylate.

2',6'-Dimethylphenyl acridine-9-thiocarboxylate (2.46 g) was suspended in 2-propanol (200 mL). Ammonium chloride (9.5 g) was added by portions to the solution followed by zinc (11.65 g). The solution was gently warmed for 2 h. TLC of the reaction mixture showed complete consumption of starting material. The solution was filtered and precipitate was washed with CH_2Cl_2 . The filtrate was concentrated to obtain a white solid which was redissolved in CH_2Cl_2 and washed with water (3 x 50 mL). The organic layer was dried over Na_2SO_4 and concentrated to yield 2.46 g of the product as a white solid. ^1H NMR (CDCl_3) δ 2.08 (s, 6H), 5.24 (s, 1H), 6.27 (br s, 1H), 6.78-7.31 (m, 11H).

c. 2',6'-Dimethylphenyl 10-methylacridanthiocarboxylate. 2',6'-Dimethylphenyl 10-methylacridan-9-thiocarboxylate (2.0 g) dissolved in 30 mL of CH_2Cl_2 under argon and

methyl triflate (6.6 mL) was added. The solution was stirred for 3 days at room temperature. The mixture was evaporated to dryness and the crude product purified by column chromatography (CH_2Cl_2) to obtain 1.85 g of the product as a white solid. ^1H NMR (CDCl_3) δ 2.09 (s, 6H), 3.46 (s, 3H), 5.09 (s, 1H), 6.97-7.36 (m, 11H).

d. 9-(2,6-Dimethylphenylthiophosphoryloxymethylidene)-10-methylacridan, bis(cyanoethyl) ester. 2,6-Dimethylphenyl 10-methylacridan-9-thiocarboxylate (700 mg) in 12 mL of dry THF was added to a solution of LDA (1.4 eq.) in 10 mL of dry THF at -78°C . The orange solution was stirred for 60 min at -78°C . A solution of POCl_3 (280 μL) and pyridine (1.57 mL) in 4 mL of THF was added dropwise. The reaction mixture was stirred for 15 min before the dry ice bath was removed. Stirring continued for 60 min. TLC (30% ethyl acetate/hexane) showed that most of the starting material was reacted.


The solution was cooled in an ice bath and treated with pyridine (1.0 mL) and 3-hydroxypropionitrile (666 μL). After stirring over night at room temperature, the precipitated pyridine-HCl was filtered and washed with THF. The combined organic solutions were evaporated in vacuo. The organic layer was dried and concentrated to obtain an orange product which was chromatographed on silica gel (ethyl acetate) to yield 80 mg of the pure product as a yellow oil. ^1H NMR (CDCl_3) δ 2.34-2.43 (m, 4H), 2.47 (s, 6H), 3.47 (s, 3H), 3.42-3.71 (m, 4H), 7.01-

8.07 (m, 11H); ^{31}P NMR (CDCl_3) δ -10.207 (p) (rel. to ext. H_3PO_4).

e. 9-(2,6-Dimethylphenylthiophosphoryloxymethylidene)-

5 10-methylacridan, disodium salt (9). A solution of the bis(cyanoethyl) phosphate compound (0.80 g) in 6 mL of acetone was purged with argon. A solution of 1M NaOH in water (293 μL) was added dropwise followed by an additional 300 μL of water and the solution stirred under
10 argon over night. The precipitate which had formed was suction filtered, washed with acetone and air-dried. The yield was 0.60 g of 9 as a white solid. ^1H NMR (D_2O) δ 2.13 (s, 6H), 3.19 (s, 3H), 6.60-8.19 (m, 11H); ^{31}P NMR (D_2O) δ 1.653 (s) (rel. to ext. H_3PO_4).

Example 10. Synthesis of Acridan Derivative 10.



10

a. 4-Fluoroacetanilide. 4-Fluoroaniline (20 g) was dissolved in 25 mL of acetic acid and cooled in an ice
25 bath. Acetic anhydride (25 mL) was added in 5 mL portions to the stirred solution. The resulting solution was poured into 200 mL of cold water and the precipitated product filtered off. The solid was washed with water and vacuum-

dried. ^1H NMR (CDCl_3) δ 2.175 (s, 3H), 7.014 (t, 2H), 7.15 (br s, 1H), 7.43-7.48 (m, 2H).

b. 4,4'-Difluorodiphenylamine. 4-Fluoroacetanilide

5 (12.0 g) was condensed with 1-bromo-4-fluorobenzene (21 mL) in the presence of 11.08 g of K_2CO_3 and 1.61 g of CuI at 190 °C for 90 h. After cooling, the mixture was filtered and the solid washed with CH_2Cl_2 . The combined organic solutions were evaporated under vacuum producing
10 ca. 20 g of a dark brown liquid. This was dissolved in 100 mL of ethanol and refluxed with 9 g of KOH for 24 h. The ethanol was evaporated and the dark residue taken up in ether and washed with water (3 x 100 mL). The ether solution was dried and concentrated and the crude product
15 purified by column chromatography on silica using 15% ethyl acetate/hexane: ^1H NMR (CDCl_3) δ 5.46 (br s, 1H), 6.947 (s, 4H), 6.97 (s, 4H); ^{13}C NMR (CDCl_3) δ 115.86, 116.16, 119.38, 119.47.

20 c. 2,7-Difluoroacridine-9-carboxylic acid. 4,4'-
Difluorodiphenylamine (5.4 g) dissolved in 50 mL of CH_2Cl_2 was added to a solution of 2.52 mL (0.11 mol) of oxalyl chloride in 30 mL of CH_2Cl_2 under an argon atmosphere at a rate to maintain reflux. The mixture was refluxed an
25 additional 35 min. The resulting solution was evaporated to remove excess oxalyl chloride. The solid was redissolved in 75 mL of CH_2Cl_2 and cooled in an ice bath. The flask was purged with argon and AlCl_3 (14.28 g) was

added in portions while the mixture was stirred. The mixture became thick during the addition and began to reflux. Reflux was maintained for an additional hour after all of the AlCl_3 had been added. The cooled mixture was
5 evaporated in vacuo and quenched with 300 mL of 3:1 ice/5 M HCl. The mixture was stirred for 1 h, the orange solid (isatin) filtered off with suction and washed with water. ^1H NMR (acetone- d_6) δ 6.92-6.97 (m, 1H), 7.35-7.61 (m, 6H).

10 The isatin product was refluxed in 110 mL of 10% KOH solution in water over night. The dark green mixture was cooled and acidified with 400 mL of 1:1 ice/5 M HCl. The solid was washed with water and air-dried yielding the acridine acid. ^1H NMR (DMSO- d_6) δ 7.81-7.94 (m, 4H), 8.31-
15 8.36 (m, 2H).

d. Phenyl 2,7-difluoroacridine-9-thiocarboxylate. 2,7-Difluoroacridine-9-carboxylic acid (1.0 g) was suspended in SOCl_2 (5 mL) and the reaction mixture was refluxed for
20 3 h. The solvent was removed under reduced pressure leaving a brown solid which was dissolved in 20 mL of CH_2Cl_2 under argon. Thiophenol (468 mg) was added followed by pyridine (2 mL). The reaction mixture was stirred overnight at room temperature. The reaction mixture was
25 diluted with CH_2Cl_2 and washed with water (3 x 50 mL). The organic layer was dried over MgSO_4 and concentrated to obtain a crude material which was chromatographed on silica gel (40 % ethyl acetate/hexane) to yield 1.2 g of

the pure product. ^1H NMR (CDCl_3) δ 7.52-7.75 (m, 9H), 8.26-8.30 (m, 2H).

e. Phenyl 2,7-difluoroacridan-9-thiocarboxylate. Phenyl

5 2,7-difluoroacridine-9-thiocarboxylate (1.2 g) was suspended in 2-propanol (125 mL) along with NH_4Cl (4.57 g). Zinc (5.5 g) was added and the reaction mixture was warmed for 2.5 h followed by a 1.5 h period at room temperature. TLC of the reaction mixture showed complete
10 conversion to a new material. The solution was filtered and the precipitate was washed with CH_2Cl_2 . The filtrate was concentrated and the light orange residue was redissolved in CH_2Cl_2 and washed with water (2 x 100 mL). The organic layer was dried over Na_2SO_4 and concentrated to
15 yield the crude product which was used without further purification. ^1H NMR (CDCl_3) δ 5.12 (s, 1H), 6.19 (br s, 1H), 6.72-7.35 (m, 11H).

f. Phenyl 2,7-difluoro-10-methylacridan-9-thio-

20 carboxylate. Phenyl 2,7-difluoroacridan-9-thiocarboxylate (1.4 g) was dissolved in CH_2Cl_2 (50 mL) under argon and methyl trifluoromethanesulfonate (5.29 g) was added. The solution was stirred overnight at room temperature. TLC showed the conversion to be ca. 50% complete so another 2
25 mL of methyl triflate was added and stirring continued for another 40 h. This reaction mixture was concentrated and the residue separated chromatographically using 20-50 % CH_2Cl_2 /hexane to obtain the product along with a minor

amount of the isopropyl ester of the acridancarboxylate as a by-product. ^1H NMR (acetone- d_6) δ 3.41 (s, 3H), 4.98 (s, 1H), 6.89-7.34 (m, 11H).

5 g. 9-(Phenylthiophosphoryloxymethylidene)-2,7-difluoro-

10-methylacridan, bis(cyanoethyl) ester. Phenyl 2,7-

difluoro-10-methylacridan-9-thiocarboxylate (500 mg, 1.3 mmol) was added to a solution of LDA (1.4 mmol) in THF -78 °C. After stirring 1 h, a solution of POCl_3 (313 mg) and

10 pyridine (1.07 g) in 4 mL of THF was added and the reaction mixture maintained at -78 °C for 1 h. The dry ice bath was removed and stirring continued for 1 h.

The solution was cooled in an ice bath and treated dropwise with pyridine and 3-hydroxypropionitrile (484 mg)

15 in 4 mL of THF. After stirring over night at room temperature, the precipitated pyridine-HCl was filtered

away and the reaction solvent evaporated in vacuo. The

residue was taken up in ethyl acetate and washed with 4 x 25 mL of water. After drying and evaporating the ethyl

20 acetate, the residue was separated chromatographically (75-80 % ethyl acetate/hexane) to yield the product was

isolated; ^1H NMR (CDCl_3) δ 2.48-2.65 (m, 4H), 3.48 (s, 3H), 4.02-4.16 (m, 4H), 6.91-7.66 (m, 11H); ^{31}P NMR (CDCl_3) δ -9.874 (p) (rel. to ext. H_3PO_4).

25


h. 9-(Phenylthiophosphoryloxymethylidene)-2,7-difluoro-

10-methylacridan, disodium salt (10). A solution of the

bis(cyanoethyl) phosphate compound (0.28 g) in 17 mL of

acetone was cooled in an ice bath and purged with argon. A solution of 1N NaOH in 1.02 mL of water was added dropwise and the solution stirred under argon for 16 h. The precipitate which had formed was suction filtered, washed with acetone and vacuum-dried. The yield was 0.236 g of 10 as an off-white solid. ^1H NMR (D_2O) δ 3.28 (s, 3H), 6.78-8.21 (m, 11H); ^{31}P NMR (D_2O) δ 1.006 (s) (rel. to ext. H_3PO_4).

10 Example 11. Synthesis of Acridan Derivative 11.



11

a. 2',2',2'-Trifluoroethyl acridine-9-thiocarboxylate.

Acridine-9-carbonyl chloride (1.89 g), made as described above was dissolved in 30 mL of CH_2Cl_2 under argon. This solution was cooled in an ice bath and 2,2,2-trifluoroethanethiol (1.0) was added dropwise followed by 3.23 g of pyridine. The reaction mixture was stirred at room temperature for 3 days. Water (20 mL) was added and the remaining precipitate filtered and discarded. The CH_2Cl_2 solution was washed with 3 x 25 mL of water, dried and evaporated to obtain the product as an off-white solid sufficiently pure to take on to the next reaction. ^1H NMR

(CDCl₃) δ 4.02-4.10 (q, 2H), 7.61-8.31 (m, 8H).

b. 2',2',2'-Trifluoroethyl acridan-9-thiocarboxylate.

2',2',2'-Trifluoroethyl acridine-9-thiocarboxylate (1.7 g)
5 was suspended in 2-propanol (150 mL) along with NH₄Cl
(7.08 g). Zinc (8.61 g) was added and the reaction mixture
was stirred for 1.5 h at room temperature. TLC of the
reaction mixture showed complete conversion to a new
material. The solution was filtered and the precipitate
10 was washed with CH₂Cl₂. The filtrate was concentrated and
the light orange residue was redissolved in CH₂Cl₂ and
washed with water (3 x 50 mL). The organic layer was dried
and concentrated to yield the crude product as an off-
white solid sufficiently pure to take on to the next
15 reaction. ¹H NMR (CDCl₃) δ 3.37-3.47 (q, 2H), 5.206 (s,
1H), 6.27 (br s, 1H), 6.78-7.27 (m, 8H).

c. 2',2',2'-Trifluoroethyl 10-methylacridan-9-thio-
carboxylate. 2',2',2'-Trifluoroethyl acridan-9-thiocarbox-
20 ylate (1.66 g) was dissolved in CH₂Cl₂ (20 mL) under argon
and methyl triflate (4.06 mL) was added. The brown
solution was stirred for 1.5 days at room temperature.
After concentrating the CH₂Cl₂, the crude product was
purified by column chromatography using 50 % CH₂Cl₂/hexane.
25 ¹H NMR (CDCl₃) δ 3.39-3.43 (q, 2H), 3.43 (s, 3H), 5.04 (s,
1H), 6.98-7.38 (m, 8H).

d. 9-(2',2',2'-Trifluoroethylthiophosphoryloxymethyl-

idene)-10-methylacridan, bis(cyanoethyl) ester. 2',2',2'-
Trifluoroethyl 10-methylacridan-9-thiocarboxylate (300 mg)
in 10 mL of THF was added to a solution of LDA (1.4 eq.)
in 10 mL of THF dropwise at -78 °C. After stirring 60 min
5 at -78 C, a solution of POCl₃ (239 mg) and pyridine (703
mg) in 4 mL of THF was added slowly. After 30 min, the dry
ice bath was removed and stirring continued for 60 min.
The solution was cooled in an ice bath and treated with :
pyridine (500 µL) and 3-hydroxypropionitrile (395 µL). A
10 brownish-yellow precipitate formed. After stirring over
night at room temperature, the precipitate was filtered
away and the reaction solvent evaporated in vacuo. The
residue was taken up in ethyl acetate, washed with 5 x 25
mL of water, dried and concentrated. The crude product was
15 separated chromatographically using CH₂Cl₂ to elute the
column and then switching solvents to a 60-100 % ethyl
acetate/hexane gradient. Two fractions were isolated, each
as mixture of two closely eluting products, the slower
eluting being the minor component. Each fraction was
20 further purified separately by prep. TLC using 25 % ethyl
acetate/hexane. The slower eluting product proved to be
the expected bis(cyanoethyl)phosphate ester. ¹H NMR
(CDCl₃) δ 2.69 (t, 4H), 3.41-3.54 (q, 2H), 3.49 (s, 3H),
4.04-4.24 (m, 4H), 7.04-7.95 (m, 8H); ³¹P NMR -8.66 (p).
25 The other product was determined by ¹H NMR, ¹³C NMR, ¹⁹F
NMR and ³¹P NMR as well as homonuclear and heteronuclear
decoupling experiments to be 9-(2',2'-difluoroethenylthio-
phosphoryloxymethylidene)-10-methylacridan,


bis(cyanoethyl) ester, derived from elimination of HF from the SCH_2CF_3 group and phosphorylation of the enolate. ^1H NMR (CDCl_3) δ 2.60-2.76 (m, 4H), 3.49 (s, 3H), 4.02-4.26 (m, 4H), 5.17 (d, 1H), 7.03-7.88 (m, 8H); ^{19}F NMR (CDCl_3) δ -76.76 (d, $J = \text{Hz}$), -78.88 (t, $J = \text{Hz}$); ^{31}P NMR (CDCl_3) δ -8.83 (p). Irradiation of the doublet at δ 5.17 did not affect any other signal in the ^1H NMR spectrum.

Irradiation of the multiplet at δ 4.02-4.26 collapsed the multiplet at δ 2.60-2.76 to a singlet and collapsed the phosphorus signal to a singlet. Irradiation of the doublet at δ 5.17 in the ^1H spectrum collapsed the triplet in the ^{19}F spectrum to a doublet.

e. 9-(2',2',2'-Trifluoroethylthiophosphoryloxymethyl)-idene)-10-methylacridan, disodium salt (11). A solution of the bis(cyanoethyl) phosphate compound (15 mg) in 4 mL of acetone was purged with argon. A solution of 1M NaOH in water (58 μL) was added dropwise followed by an additional 100 μL of water and the solution stirred under argon over night. TLC showed complete deprotection. The mixture was concentrated to dryness, producing an off-white solid which was washed with 20 % CH_2Cl_2 /acetone and air-dried. ^1H NMR (D_2O) δ 3.399 (s, 3H), 3.45-3.61 (m, 2H), 7.04-8.18 (m, 8H); ^{31}P NMR (D_2O) δ 1.465 (s) (rel. to ext. H_3PO_4).

Example 12. Synthesis of Acridan Derivative 12.

5



12

a. 4'-Chlorophenyl acridine-9-thiocarboxylate. 4-

10 Chloro-thiophenol (4.75 g) and 7.22 g of acridine-9-carbonyl chloride were dissolved in 100 mL of CH_2Cl_2 , followed by 12.1 mL of pyridine. The reaction mixture became orange-brown. The mixture was stirred overnight at room temperature under argon. After evaporation of

15 solvent, the solids were washed with 100 mL of hexanes, filtered, washed with another 100 mL of hexanes, filtered, and then with a 500 mL of water, filtered and air-dried. The thioester was obtained (8.71 g) as a slightly brownish yellow solid. ^1H NMR (CDCl_3) δ 7.47-7.50 (m, 2H), 7.58-7.67 (m, 4H), 7.81-7.86 (m, 2H), 8.12 (d, 2H), 8.29

20 (d, 2H).

b. 4'-Chlorophenyl acridan-9-thiocarboxylate. 4'-

25 Chlorophenyl acridine-9-thiocarboxylate (2.0 g) was dissolved in CH_2Cl_2 (25 mL). The solution was purged with argon and then 3.72 g of zinc powder added followed by 0.45 mL of acetic acid. TLC showed the starting material was consumed in 20 min. The mixture was filtered and the

solids washed with CH_2Cl_2 . The combined CH_2Cl_2 solutions were washed with water (3 x 50 mL) and dried. A small amount of the product was purified by prep. TLC for analytical characterization. the remainder of the product was used without further purification. ^1H NMR (CDCl_3) δ 5.22 (s, 1H), 6.28 (s, 1H), 6.79-7.31 (m, 12H).

c. 4'-Chlorophenyl 10-methylacridan-9-thiocarboxylate. 4'-Chlorophenyl acridan-9-thiocarboxylate (2.0 g) was dissolved in CH_2Cl_2 (30 mL) under argon and methyl triflate (6.5 g) was added. The brown solution was evaporated and the crude product purified by column chromatography using 30 % ethyl acetate/hexanes. The pure product (1.8 g) was thereby obtained. ^1H NMR (CDCl_3) δ 3.47 (s, 3H), 5.09 (s, 1H), 7.02-7.39 (m, 12H).


d. 9-(4-Chlorophenylthiophosphoryloxymethylidene)-10-methylacridan, bis(cyanoethyl) ester. 4-Chlorophenyl 10-methylacridan-9-thiocarboxylate (0.70 g) in 10 mL of anhydrous THF was added dropwise to a solution of LDA (1.4 eq.) in THF at -78 C. After stirring 60 min at -78 C, the yellow solution was treated with a solution of POCl_3 (0.517 g) and pyridine (1.52 mL) in 4 mL of THF slowly. The dry ice bath was removed after 30 min and stirring continued for 1 h. The solution became yellow and formed a precipitate.

The mixture was cooled in an ice bath and treated with 3-hydroxypropionitrile (0.89 g) and 1.0 mL of pyridine.

The ice bath removed after the addition was complete. After stirring over night at room temperature, the precipitated pyridine-HCl was filtered off and washed with THF. The combined filtrates were evaporated in vacuo and the brown material obtained was dissolved in ethyl acetate and washed with 4 x 25 mL of water. The ethyl acetate solution was dried and concentrated. The residue was separated by column chromatography using 80-100 % ethyl acetate/hexanes from which the 0.325 g of the product was isolated; ^1H NMR (CDCl_3) δ 2.48-2.64 (m, 4H), 3.53 (s, 3H), 3.86-4.16 (m, 4H), 6.94-7.94 (m, 12H); ^{31}P NMR (D_2O) δ -9.48 (p) (rel. to ext. H_3PO_4).

e. 9-(4-Chlorophenylthiophosphoryloxymethylidene)-10-methylacridan, disodium salt (12). A solution of the bis(cyanoethyl) phosphate compound (0.325 g) in 10 mL of acetone was purged with argon. A solution of 2.5 M NaOH (473 μL) was added followed by an additional 500 μL of water. The solution was stirred under argon over night. The precipitate which had formed was suction filtered and air-dried. The mother liquor was found to contain the mono(cyanoethyl)-protected compound (140 mg). A second crop of the disodium salt was obtained by repeating the NaOH deprotection. ^1H NMR (D_2O) δ 3.35 (s, 3H), 6.92-7.36 (m, 10H), 7.78 (d, 1H), 8.20 (d, 1H); ^{31}P NMR (D_2O) δ 1.22 (s) (rel. to ext. H_3PO_4).

Example 13. Synthesis of Acridan Derivative 13.



5

13

a. Naphthyl acridine-9-thiocarboxylate. 2-Naphthal-
 enethiol (48.81 g) and the acridine-9-carbonyl chloride
 10 prepared from 65.17 g of acridine-9-carboxylic acid were
 dissolved in 100 mL of CH_2Cl_2 followed by the addition of
 120 mL of pyridine. The mixture was stirred overnight at
 room temperature under argon. After evaporation of solvent,
 the solids were washed with 500 mL of hexanes, filtered,
 15 washed with another 500 mL of hexanes, filtered, and then
 washed with 600 mL of water, filtered and air-dried
 overnight. The thioester was dissolved in 1500 mL of
 CH_2Cl_2 , dried over sodium sulfate, filtered, and dried in
 vacuo. The thioester was obtained (94.67 g) as a brown
 20 solid. ^1H NMR (CDCl_3) δ 7.54-8.00 (m, 11H), 8.17-8.31 (m,
 4H).

b. Naphthyl 10-methylacridinium-9-thiocarboxylate
triflate. Naphthyl acridine-9-thiocarboxylate (26.38 g) was
 25 suspended in CH_2Cl_2 (200 mL). Methyl
 trifluoromethanesulfonate (24.5 mL) was added and the
 mixture left to stir overnight. The mixture was filtered
 and the solids washed with CH_2Cl_2 (300 mL) and hexanes (500

ml). After air drying, the product (28.83 g) was obtained as a yellow solid. ¹H NMR (acetone-d₆) δ 5.20 (s, 3H), 7.66-7.75 (m, 2H), 7.88-7.92 (m, 1H), 8.03-8.09 (m, 2H), 8.15 (d, 1H), 8.26 (t, 2H), 8.48 (s, 1H), 8.61-8.68 (m, 2H), 8.80 (d, 2H), 9.03 (d, 2H).

c. Naphthyl 10-methylacridan-9-thiocarboxylate. Naphthyl 10-methylacridinium-9-thiocarboxylate triflate (65.50 g) was suspended in CH₂Cl₂ (1000 mL) under argon and glacial acetic acid (21.2 ml) and zinc (40.43 g) were added. After stirring overnight, TLC showed the disappearance of the starting material and the formation of a new product. The reaction mixture was filtered through a bed of silica gel and the CH₂Cl₂ removed in vacuo. The slightly yellow solid obtained was stirred in isopropanol (500 ml), filtered, washed with 500 ml more isopropanol, and allowed to air dry. The pure product (46.36 g) was thereby obtained. ¹H NMR (CDCl₃) δ 3.47 (s, 3H), 5.13 (s, 1H), 7.00-7.06 (m, 4H), 7.25-7.49 (m, 7H), 7.70-7.79 (m, 4H).

20

d. 9-(Naphthylthiophosphoryloxymethylidene)-10-methylacridan, bis(cyanoethyl) ester. Naphthyl 10-methylacridan-9-thiocarboxylate (17.32 g) in 600 mL of anhydrous THF was added dropwise to a solution of LDA (1.25 eq.) in THF at -78 °C. After stirring 90 min at -78 °C, the orangish brown solution was treated with a solution of POCl₃ (20.80 g) and pyridine (50 mL) in 150 mL of THF slowly. The dry ice bath was removed after 60 min and

25

stirring continued for 1 h. The solution became brown and formed a precipitate.

The mixture was treated with 3-hydroxypropionitrile (27.8 ml). After stirring over night at room temperature, the precipitated pyridine-HCl was filtered off and washed with THF. The combined filtrates were evaporated in vacuo and the brown oil obtained was separated by column chromatography using 50-100 % ethyl acetate/hexanes from which the product was isolated. The yellow oil was dissolved in CH_2Cl_2 (300 ml), washed with water (450 ml), dried over sodium sulfate, filtered, and dried in vacuo. This yielded the product (17.76 g) as a yellow solid. ^1H NMR (CDCl_3) δ 2.33-2.53 (m, 4H), 3.53 (s, 3H), 3.82-4.06 (m, 4H), 6.93 (t, 1H), 7.03 (d, 1H), 7.10-7.18 (m, 2H), 7.25-7.55 (m, 5H), 7.79-7.92 (m, 5H), 8.02 (d, 1H); ^{31}P NMR (CDCl_3) δ -9.69 (p) (rel. to ext. H_3PO_4).

e. 9-(Naphthylthiophosphoryloxymethylidene)-10-methylacridan, disodium salt (13). A solution of the bis(cyanoethyl) phosphate compound (17.28 g) in 200 mL of acetone was purged with argon. A solution of NaOH (2.76 g) in 50 ml of water was added and the solution was stirred under argon over night. The precipitate which had formed was suction filtered washed with 20 % water in acetone (300 ml) and dried under vacuum. The product (15.07 g) was obtained as a light yellow solid. ^1H NMR (D_2O) δ 3.22 (s, 3H), 6.67 (d, 1H), 6.87 (t, 1H), 7.01 (t, 1H), 7.08-7.15 (m, 2H), 7.23 (d, 1H), 7.31-7.44 (m, 3H), 7.51 (s, 1H), 7.59

(d, 2H), 7.73 (d, 1H), 7.86 (d, 1H), 8.25 (d, 1H); ^{31}P NMR (D_2O) δ 1.30 (s) (rel. to ext. H_3PO_4).

Example 14. Chemiluminescent Detection of Alkaline

5 Phosphatase with Compound 1. An effective reagent composition for chemiluminescent detection of alkaline phosphatase comprised 0.1 M tris buffer, pH 8.5, 0.88 mM magnesium salt, 0.33 mM acridan phosphate 1 (from a 1:100 dilution of a 0.033 M methanol solution) and 0.01 mg/mL of
10 the Enhancer A (polyvinylbenzyltributylphosphonium chloride co-polyvinylbenzyltrioctylphosphonium chloride, containing a 3:1 ratio of tributyl:trioctyl groups). Reaction of 100 μL of this composition with 10^{-13} mol of AP at 25 °C in a test tube housed in a Turner TD-20e luminometer produced
15 easily measurable blue chemiluminescence which reached maximum intensity in 2 min.

Example 15. Chemiluminescent Detection with Compounds 2-13.

In the manner of Example 14, compositions containing each
20 of compounds 2-13 were reacted with AP at 25 °C. Each produced easily measurable chemiluminescence discernable above the background measured in the absence of AP.

It will be obvious to one skilled in the art that other compounds of formula I in addition to those specifically
25 illustrated herein will function in like manner in the present methods.

Example 16. Chemiluminescent Detection of Alkaline

Phosphatase with Compound 5. An effective reagent composition for chemiluminescent detection of alkaline phosphatase comprised 0.2 M 2-methyl-2-amino-1-propanol buffer, pH 9.6, 0.88 mM Mg^{2+} , 0.66 mM acridan phosphate 5 (from a 1:100 dilution of a 0.033 M methanol solution) and 0.5 mg/mL of the Enhancer A. Reaction of 100 μ L of this composition with 10^{-13} mol of AP at 25 °C in a test tube housed in a Turner TD-20e luminometer produced easily measurable blue chemiluminescence which reached maximum 10 intensity in 2 min. Light intensity measured between 1 and 10 min correlated with amount of enzyme in the range 8×10^{-14} mol to 8×10^{-18} mol.

Example 17. Kinetic Profile of Chemiluminescence Intensity.

15 The rapid detection speed afforded by compositions containing the acridan phosphate 1 is shown by the plot in Figure 1. Reaction of 100 μ L of the reagent composition of Example 14 with 8×10^{-16} mol of AP at 25 °C caused a burst of light emission which achieved maximum intensity in ca. 2 20 min. The figure also shows for comparison the chemiluminescence profile of the ester phenyl 10-methylacridan-9-carboxylate in a reagent composition similar to that described in Example 14 but containing the ester in place of 1. Addition of AP to the latter solution 25 of the ester produced no effect. The much lower light intensity and slower rise time from spontaneous autoxidation of the ester is incompatible with a scheme in which emission from reaction of 1 is simply due to

generation of the ester (via removal of the phosphate protecting group from the enol phosphate) by AP followed by autoxidation of the ester.

5 Example 18. Spectroscopic Study of Reaction of 1 with AP.

The progress of the reaction between 3 mL of the reagent composition of Example 14 and 2.4×10^{-12} mol of AP at ambient temperature was monitored by UV-visible spectroscopy with a diode array detector. The spectrum was
10 acquired at 5 min intervals. The evolution of the spectrum revealed in Figure 2 showed a complex pattern consisting of multiple intermediates and products. Curves labeled A-F represent scans taken 0, 5, 10, 20, 40 and 60 min after addition of enzyme.

15

Example 19. Chemiluminescence Spectrum. A chemiluminescence spectrum was obtained using a device constructed by Dr. Barry Schoenfelner of Lumigen. The reagent composition of Example 14 (200 μ L) was reacted with AP (8×10^{-13} mol) in a
20 sample holder consisting of a test tube inside a light tight box. The emitted light was collected by a lens, and dispersed by a grating monochromator onto a CCD camera element. No correction for the change in intensity with time is required since this system images all wavelengths
25 of visible light simultaneously. The spectrum of light emitted, shown in Figure 3, reveals a broad emission with a maximum intensity at about 430 nm. This spectrum matches the fluorescence of N-methylacridone. The apparent fine

structure in the spectrum is the result of instrument noise.

Example 20. Effect of pH on Chemiluminescent Reaction of 1
with Alkaline Phosphatase. Chemiluminescent detection of
 5 alkaline phosphatase with acridans of the present invention
 can be performed over a broad range of pH. Solutions
 comprising a 0.1 M buffer (either tris or 2-methyl-2-amino-
 1-propanol, 221), pH 7-10.5, 0.88 mM Mg^{2+} , 0.33 mM acridan
 10 phosphate 1 (from a 1:100 dilution of a 0.033 M methanol
 solution) and 0.01 mg/mL of Enhancer A were prepared and
 the background light emission measured after 10 min at 25
 °C. An aliquot of an AP solution containing 8×10^{-16} mol of
 enzyme was injected and light intensity monitored until
 15 maximum intensity was reached. S/B represents the ratio of
 maximum light intensity to the light intensity in the
 absence of enzyme.

Table 2.

	<u>Entry</u>	<u>pH</u>	<u>Buffer</u>	<u>S/B</u>
20	1	7.0	tris	37
	2	7.5	tris	144
	3	8.0	tris	273
	4	8.5	tris	280
	5	9.0	tris	195
25	6	9.0	221	200
	7	9.5	221	87
	8	10.0	221	13
	9	10.5	221	1.3

Example 21. Enhancement of Chemiluminescence by

Surfactants. The enhancement of chemiluminescence, defined as the increase in light intensity over the level in the absence of the test substance, was assessed under a

5 standard set of test conditions in the reaction of acridan
1 with AP in the presence of various surfactant compounds.
The concentration of each surfactant producing the greatest
enhancement was determined independently. The value in the
column marked S/B represents the ratio of light intensities
10 with 10^{-16} mol of AP and that with no enzyme. The column
marked Rel. presents the same ratio normalized to the
results obtained with no enhancer (entry 11).

Table 3

	<u>Entry</u>	<u>Surfactant</u>	<u>Concentration</u>	<u>S/B</u>	<u>Rel.</u>
	1	A	0.01 g/L	562	40.1
	2	B	0.01	431	30.8
5	3	C	0.01	389	27.8
	4	D	0.25	295	21.1
	5	E	0.10	162	11.6
	6	F	0.05	120	8.6
	7	G	0.10	63	4.5
10	8	H	0.10	26	1.9 *
	9	I	0.43	24	1.7
	10	J	0.10	17	1.2
	11	None	----	14	1

* Measured at 2 min.

15 A: poly(vinylbenzyltributylphosphonium chloride)-co-poly-(vinylbenzyltrioctylphosphonium chloride) (3:1 ratio of tributyl:trioctyl groups)

B: polyvinylbenzylbenzyldimethylammonium chloride

C: poly(vinylbenzyltributylphosphonium chloride)-co-poly-(vinylbenzyltrioctylphosphonium chloride)-co-poly(vinylbenzylfluorescein) (75:25:1)

D: poly(vinylbenzyltributylphosphonium chloride)

E: 1-trioctylphosphoniummethyl-4-tributylphosphonium-methylbenzene dichloride

25 F: Tween™ 20, polyoxyethylene(20) sorbitan monolaurate

G: 1-trioctylphosphoniummethyl-4-tributylammonium-methylbenzene, mixed bromide chloride salt

H: SDS, sodium dodecyl sulfate

I: CTAB, cetyltrimethylammonium bromide

J: 1-trioctylammoniummethyl-4-tributylammoniummethylbenzene
dichloride

The S/B values in entries 4-11 do not represent the maximum
5 enhancement obtainable since light intensity continues to
rise for > 1 hour when these enhancers are used.

Example 22. Effect of the Concentration of Enhancer A on
Light Intensity and Kinetics with Acridan Phosphate 1. A

10 concentration dependence study was conducted in order to
determine the amount of Enhancer A which produced the
highest signal/background (S/B) level in the
chemiluminescent reaction of acridan phosphate 1 with AP.
Solutions of 1 in 0.1 M tris buffer, pH 8.5 containing 0.88
15 mM Mg^{+2} and 0.5, 0.25, 0.1, 0.05 or 0.025 mg/mL of Enhancer
A were prepared. Aliquots (100 μ L) were equilibrated at
25 °C and reacted with 8×10^{-16} moles of AP. The S/B values
reported in Table 4 show that maximum light enhancement
occurs with 0.005-0.01 g/L of enhancer.

20 Table 4.

	<u>Entry</u>	<u>Concentration</u>	<u>S/B 2 min</u>	<u>S/B 10 min</u>
	1	0.0025 g/L	431	262
	2	0.005	714	500
	3	0.010	660	562
25	4	0.025	399	360
	5	0.050	247	240
	6	0.10	212	163
	7	0.50	40	84

Example 23. Linearity and Sensitivity of Detection of Alkaline Phosphatase with Acridan Phosphate 1. The

sensitivity and linearity of detection of AP was determined using a reagent composition of the present invention
5 containing acridan phosphate 1. To each of 3 wells in a 96-well white microplate 10 μ L dilutions of AP containing between 8×10^{-15} mol and 8×10^{-20} mol of enzyme were added. A 100 μ L aliquot of detection reagent comprising 0.33 mM solution acridan phosphate 1 in 0.1 M tris buffer,
10 pH 8.5, 0.88 mM $MgCl_2$, 0.01 mg/mL of Enhancer A and 1 % methanol was injected. Light intensities were measured after 2.5 min. Figure 4 shows the linear detection of alkaline phosphatase. The term S-B refers to the chemiluminescence signal (S) in RLU in the presence of AP
15 corrected for background chemiluminescence (B) in the absence of AP. The calculated detection limit (twice the standard deviation of the background) was determined to be 1.2×10^{-19} mol under these conditions. A similar result was obtained using light intensities measured after 10 min.

20

Example 24. Chemiluminescent Detection of AP by pH Change.

An alternate way to detect the light emitted by reaction of AP with 1 is to incubate a reagent composition containing 1 with AP at a first pH in the absence of enhancer and then
25 rapidly raise the pH by injecting a solution of a strong base containing enhancer to cause a burst of light emission. A set of conditions which permitted the detection of $< 10^{-18}$ mol of AP was devised.

A reagent composition (100 μ L) comprising 0.33 mM acridan phosphate 1 in 0.1 M tris buffer, pH 9, 0.88 mM $MgCl_2$, was reacted with 10 μ L of AP (8×10^{-15} - 8×10^{-20} mol) or 10 μ L of water as a reagent blank for 4 min at ambient temperature in the wells of a black microplate. A solution containing 0.5 mg/mL of Enhancer A in 1 N NaOH (100 μ L) was injected and light intensity integrated for 10 sec. Light intensity was a linear function of the amount of AP between 8×10^{-15} and 8×10^{-19} mol of AP as shown in Figure 5.

Example 25. Chemiluminescent Detection of Acid Phosphatase and Inhibition by Tartrate. A reagent composition for chemiluminescent detection of acid phosphatase (AcP) comprised 0.1 M tris buffer, pH 7, 0.33 mM acridan phosphate 1 (from a 1:100 dilution of a 0.033 M methanol solution) and 0.01-0.1% of Enhancer A. Reaction of 100 μ L of this composition with AcP (Sigma AcP LIN-TROL, reconstituted to 2.0 mL, 83 U/L) at ambient temperature in a test tube housed in a Turner TD-20e luminometer produced chemiluminescence which reached maximum intensity nearly instantaneously and remained nearly constant for several minutes. Addition of the protein bovine serum albumin to the detection reagent at 0.1% prolonged the duration of light emission significantly.

Addition of 5 μ L of 0.04 M tartrate in citrate buffer (0.09 M, pH 4.8) caused complete extinction of light emission due to inhibition of enzyme activity by tartrate,

a specific inhibitor for prostatic acid phosphatase. The quenching of emission was not due to a reduction of the pH since addition of 10 μ L of citrate buffer caused only a 25% reduction in light intensity.

5

Example 26. Chemiluminescent Detection of AcP by pH Change.

An alternate way to detect the light emitted by reaction of AcP with 1 is to incubate a reagent composition containing 1 and enhancer with AcP at a first pH and then rapidly
10 raise the pH by injecting a solution of a strong base to cause a burst of light emission. A reagent composition (100 μ L) comprising 0.33 mM acridan phosphate 1 in 0.1 M tris buffer, pH 7.0, 0.1% BSA and 0.01% Enhancer A was reacted with 3 μ L of AcP (Sigma AcP LIN-TROL, reconstituted to 2.0
15 mL, 83 U/L) for 8 min at ambient temperature in a test tube. A solution of 1 N NaOH (50 μ L) was injected and light intensity integrated for 4 min. Light intensity was three times the reagent blank.

20 Example 27. Western Blot Assay using PVDF Membrane.

Compositions of the present invention were used to detect and quantify a protein, human transferrin in a Western blot with an AP-labeled antibody on a polyvinylidene difluoride (PVDF) membrane. Dilutions of transferrin containing from
25 5000, 1000, 180, 30 and 5 pg, respectively, of protein were electrophoresed and transferred to PVDF membranes (Millipore, Bedford, MA). The transferrin bands were blocked with and then reacted sequentially with goat anti-

human transferrin and rabbit anti-goat-AP conjugate. The membranes were soaked briefly with reagents comprising 0.2 M 221 buffer, pH 9.6 containing 0.88 mM MgCl_2 , 0.66 mM acridan phosphate 1 and 0.1 to 1 mg/mL of Enhancer A. The membranes were placed between transparent plastic sheets and exposed to X-ray film. Figure 6 demonstrates the detection of human transferrin after 15 min with a 1 min exposure using a reagent composition containing 0.1 mg/mL of enhancer. The light produced using these compositions with varying amounts of the enhancer led to intense emission which could be imaged for at least 2 weeks.

Surprisingly it has been found that compositions containing 0.5 mg/mL of Enhancer A produced light emission whose intensity was nearly constant for 4 days with no change in the relative band intensities. Light emission produced on PVDF membranes persisted at useful levels for very long periods of time, exceeding a month. The 5 pg band could still be imaged with only a 20 min exposure after 5 weeks. This duration of emission at useful levels is without precedent.

Example 28. Western Blot using Nitrocellulose Membrane. A Western blot assay according to the procedures in Example 27 was performed using nitrocellulose membrane as the solid phase. Transferrin standards in the range of 5000 - 30 pg were used. A detection reagent which permitted detection of all levels of protein comprised 0.2 M 22 buffer, pH 9.6, 0.88 mM MgCl_2 , 0.33 mM acridan phosphate 1 and 0.5 mg/mL of

Enhancer A. Detection could be performed over several hours.

Example 29. Southern Blot Assay with Chemiluminescent

5 Detection. The use of a composition of the present invention for the chemiluminescent detection of DNA on nitrocellulose membranes by the technique of Southern blotting is demonstrated in the following example.

Avidin-AP conjugate was obtained from Cappel Products
10 (Durham, NC). Bovine serum albumin (heat shocked) was purchased from Sigma Chemical Co. (St. Louis, MO). Human genomic DNA and human transferrin receptor cDNA were from Clontech (Palo Alto, CA). Biotinylated lambda DNA/HindIII fragments, biotin-7-dATP and the Nick Translation kit were
15 from Life Technologies, restriction endonuclease HindIII was from Boehringer-Mannheim (Indianapolis, IN), and nitrocellulose membrane from Schleicher & Schuell Inc (Keene, NH). X-ray film was X-OMAT AR from Kodak (Rochester, NY).

20 Human genomic DNA (19.5 µg) was cleaved with HindIII to completion and divided into 13 µg and 6.5 µg portions. Restricted DNA was purified by phenol/chloroform extraction and ethanol precipitation. Purified DNA was separated by electrophoresis on 0.75% agarose gel with 40 mmol/L Tris-
25 acetate, 2 mmol/L EDTA, pH 8.0 as the elution buffer. Following electrophoresis, the gel was rinsed with water, equilibrated for 12 min with 0.25 mol/L HCl, rinsed with water again, incubated in 0.5 mol/L NaOH, 1.5 mol/L NaCl-

for 15 min then in a fresh change of the same solution for 30 min, rinsed with water, and incubated in three changes of 1 mol/L tris, 1.5 mol/L NaCl, pH 7.5 for 15 min each.

Nitrocellulose membrane was soaked sequentially with
5 water for two min and 10X SSC for 30 min. DNA was transferred to the membrane by capillary blotting overnight in 10X SSC. The membrane was washed by gentle agitation in 10X SSC for 10 min at room temperature and air-dried on Whatman 3MM blotting paper for 30 min and baked at 80° C
10 under vacuum for 2 h.

The membrane was soaked in 6X SSPE (20X SSPE is 3 mol/L NaCl, 0.2 mol/L NaH_2PO_4 , 20 mmol/L EDTA, pH 7.4) followed by prehybridization for 3 h at 42° C in pre-hybridization solution: 6X SSPE, 50% freshly deionized formamide,
15 filtered 5X Denhardt's solution (50X is 1% Ficoll 400, 1% PVP, 1% BSA (initial fraction by heat shock)), filtered 1% SDS and 200 µg/mL sheared, denatured herring sperm DNA. The hybridization probe, human transferrin receptor cDNA was labeled with biotin-7-dATP by nick translation according to
20 the manufacturers instructions. Genomic DNA was hybridized overnight at 42° C in 6X SSPE, 45% formamide, filtered 5X Denhardt's, 1% SDS, 200 µg/mL herring sperm DNA with 300 µg/mL denatured biotinylated probe. The biotinylated probe was denatured by boiling 4 min and cooling 10 min at 0° C.
25 The membrane was washed twice with 0.5X SSC, 0.4% SDS at 25° C for 5 min, three times with 0.5X SSC, 0.4% SDS at 55° C for 10 min, once with 2X SSC at 25° C for 5 min, twice with TBS (50 mmol/L Tris-HCl, pH 7.4, 0.15 mol/L NaCl) for

3 min. After the wash step, the membrane was blocked at 63° C for 1 h in filtered 3% BSA, 100 mmol/L Tris-HCl, pH 7.4, 0.15 mol/L NaCl and washed in T-TBS (0.05% Tween 20 in TBS) for 1 min.

5 The blocked membrane was incubated with a 1:2000 dilution of avidin-AP in T-TBS for 12 min followed by four fresh changes of T-TBS for 5, 10, 15 and 20 min followed by a final wash with TBS for 5 min. Excess buffer was drained off and blots soaked in the detection reagent described in
10 Example 27 for 3 min. Excess reagent was drained off, and the blots placed between transparent sheets and exposed to X-ray film.

 The single copy gene was detected at 10.5 and 5.2 kbp by soaking the membrane with detection reagent and exposure
15 to x-ray film for varying lengths of time. After an incubation period of 12 min, a band corresponding to the single copy gene in both fractions was visible with a 5 min exposure to film using a reagent of the present invention. Shorter incubation times also produced excellent images.
20 Multiple exposures could be easily performed for a day.

Example 30. Effect of Lucigenin on the Chemiluminescence Profile. The higher light intensity afforded by the addition of lucigenin to the reaction of acridan phosphate
25 5 with AP is shown by the plots in Figure 7. Reaction of 100 µL portions of a solution of 5 (0.66 mM) in 0.1 M tris buffer, pH 8.8 containing 0.88 mM MgCl₂, with 8×10^{-16} mol of AP at 25 °C were conducted in the presence and absence.

of 6.4 μM lucigenin. Figure 7 shows the much higher light intensity resulting from incorporation of lucigenin into the solution.

- 5 Example 31. Chemiluminescent Detection of Alkaline Phosphatase with Compound 5. A highly effective reagent composition for chemiluminescent detection of alkaline phosphatase comprised 0.1 M tris buffer, pH 8.8, 6.4 μM lucigenin, 0.66 mM compound 5, 1 mg/mL SDS, 0.01 mg/mL
- 10 Na_2SO_3 , 0.033% (w/v) TWEEN 20 and 0.88 mM MgCl_2 . Reaction of 100 μL of this composition with 8×10^{-16} mol of AP at 25 °C produced chemiluminescence which reached maximum intensity in 2 min. Figure 8 shows the comparison with a reagent composition containing 5 and the cationic polymeric
- 15 surfactant enhancer poly(vinylbenzyltributylphosphonium chloride)-co-poly-(vinylbenzyltrioctylphosphonium chloride), containing about a 3:1 ratio of tributyl:trioctyl groups (0.25TB/TO), triggered at 37 °C. Compositions containing the latter polymeric surfactant were found
- 20 previously to be the highest levels of chemiluminescence in the absence of a CAC.

- Example 32. Linearity and Sensitivity of Detection of Alkaline Phosphatase. The sensitivity and linearity of
- 25 detection of AP was determined using a reagent composition containing compound 5 and lucigenin (reagent A) and compared to the results achievable using a reagent composition containing compound 5 and 0.25TB/TO (reagent

B). The reagents have the following composition:

	<u>Reagent A</u>	<u>Reagent B</u>
	Compound 5, 0.66 mM	Compound 5, 0.66 mM
	0.1 M tris buffer, pH 8.8	0.2 M 221 buffer, pH 9.6
5	0.01 or 0.88 mM MgCl ₂	0.88 mM MgCl ₂
	lucigenin, 6.4 μM	0.25TB/TO, 0.5 mg/mL
	SDS, 1 mg/mL	
	Na ₂ SO ₃ , 0.01 mg/mL	
	TWEEN 20, 0.033% (w/v)	

- 10 To each of 3 wells in a 96-well white microplate, 10 μL dilutions of AP containing between 8×10^{-13} mol and 8×10^{-22} mol of enzyme were added 100 μL aliquots of detection reagent A or B. Light intensities were measured after 2.5 min. Figure 9 shows the linear detection of alkaline
- 15 phosphatase. The calculated detection limit (two standard deviations over the background) was determined to be 3.5×10^{-21} mol for reagent A and 1.2×10^{-19} mol for reagent B.

Example 33. Chemiluminescent Detection with Compounds 1-

- 20 13. In the manner of Example 30, compositions containing 0.66 mM of each of compounds 1-13 with 0 and 6.4 μM lucigenin were reacted with 8×10^{-16} moles of AP at 25 °C. Each demonstrated significantly more intense chemiluminescence when lucigenin was in the reaction
- 25 solution.

Example 34. Linearity and Sensitivity of Detection of AP using Compounds 7-12. In the manner of Example 32,

reagents were prepared according to the composition of reagent A substituting 0.66 mM of each of compounds 7-12 in place of compound 5. To 100 μ L portions of each of these reagents was added 10 μ L dilutions of AP containing between 8×10^{-16} mol and 8×10^{-22} mol of enzyme or 10 μ L of water for the reagent blank. In each case, the reaction solution containing 8×10^{-21} mol of enzyme produced a peak signal within about 2 minutes which was higher than the blank. Figures 10 and 11 and 18 depict the results for compounds 7, 8, and 12 respectively.

Example 35. Time profile of Chemiluminescence using Compound 7. The rapid generation of chemiluminescence emitted by a reagent composition containing compound 7 is demonstrated in the following experiment. A 100 μ L portion of a reagent consisting of 0.1 M tris buffer, pH 8.8, 6.4 μ M lucigenin, 0.66 mM acridan phosphate 7), 1 mg/mL sodium dodecyl sulfate, 0.01 mg/mL Na_2SO_3 , 0.033% (w/v) TWEEN 20 and 0.88 mM MgCl_2 was reacted with 8×10^{-16} mol of AP at room temperature. The chemiluminescence intensity reached a stable plateau in under 5 sec as shown in Figure 12.

Example 36. Increase in Chemiluminescence Intensity with Various CACs. Test solutions were prepared containing 21 mM of CAC 1-11 in DMSO. These were diluted 1:1000 into a solution of acridan phosphate compound 5 in 0.1 M tris buffer, pH 8.8 containing 0.88 mM MgCl_2 . Triplicate 100 μ L portions of each of the final reagent compositions were

reacted with 4×10^{-16} mol of AP at room temperature in a white microwell plate. Table 5 shows the effect on plateau intensity (S), measured at 36 min and signal/background (S/B) for each CAC. Each produced higher peak signal S and S/B than a control without CAC.

Table 5.

	<u>CAC</u>	<u>S at 36 min</u>
	1	2602
	2	45.3
10	3	5963
	4	4645
	5	34.7
	6	72.3
	7	61.6
15	8	19.1
	9	781
	10	21.1
	11	9.3
	None	2.8
20	1: Lucigenin	
	2: Basic Red 29	
	3: Basic Blue 66	
	4: Basic Blue 41	
	5: 3,3'-Diethylthiadibocyanine iodide	
25	6: 3,3'-Diethyl-9-methylthiacarbocyanine iodide	
	7: 3,3'-Diethylselenacarbocyanine iodide	
	8: 3,3'-Diethylthiacyanine iodide	
	9: IR-1040	

10: 5-[3-Ethoxy-4-(3-ethyl-5-methyl-2(3H)-benzothiazolyl-
idene)-2-butenylidene]-3-ethyl-2-[(3-ethyl-4,5-
diphenyl-2(3H)-thiazolylidene)methyl]-4,5-dihydro-4-
oxothiazolium iodide

5 11: IR-786 perchlorate

Example 37. Effect of Concentration of CAC on

Chemiluminescent Reaction of 5 with AP. Chemiluminescent
detection of AP with compositions of the present invention
10 can be performed over a broad range of concentration of
CAC. Compositions containing 0.66 mM compound 5 in 0.1 M
tris buffer, pH 8.8 containing 0.88 mM MgCl₂, were prepared
with varying concentrations of lucigenin ranging from 6.4
mM to 6.4 nM by 10-fold serial dilutions of a stock
15 lucigenin solution. The control contained no lucigenin.
Test solutions (100 µL, 5 replicates) containing between
6.4 mM and 6.4 nM reacted with 4×10^{-16} mol of AP produced
a higher signal (S) than the control after a 3.5 min
incubation at room temperature.

20

Example 38. Effect of Concentration of Anionic Surfactant
on Chemiluminescent Reaction of 5 with AP.

Chemiluminescent detection of AP with compositions of the
present invention can be performed over a broad range of
25 concentration of the anionic surfactant SDS. Compositions
containing 0.66 mM compound 5 and 0.88 mM MgCl₂ in 0.1 M
tris buffer, pH 8.8 with varying concentrations of SDS
ranging from 10 mg/mL to 10 µg/mL were prepared. The

control contained no SDS. Test solutions (100 μ L, 5 replicates) reacted with 4×10^{-16} mol of AP produced a faster rise to peak signal at room temperature than the control.

5

Example 39. Effect of Concentration of Nonionic Surfactant on Chemiluminescent Reaction of 5 with AP.

Chemiluminescent detection of AP with compositions of the present invention can be performed over a broad range of concentration of the non-ionic surfactant TWEEN 20. Compositions according to Reagent A in Example 32 were prepared but with 21 μ M lucigenin and varying concentrations of the non-ionic surfactant ranging from 1.0 % to 0.001 % and lacking the Na_2SO_3 . The control contained no non-ionic surfactant. The test solutions (100 μ L, 5 replicates) and the controls were reacted with 4×10^{-16} mol of AP at room temperature. Each test solution was effective in extending the duration of chemiluminescence at or near maximum intensity.

20

Example 40. Effect of Concentration of Sodium Sulfite on Chemiluminescent Reaction of 5 with AP.

Chemiluminescent detection of AP with compositions of the present invention can be performed over a broad range of concentration of sodium sulfite (Na_2SO_3). Compositions according to Reagent A in Example 32 were prepared but with varying concentrations of Na_2SO_3 ranging from 1.0 mg/mL to 1.0 μ g/mL. The control contained no Na_2SO_3 . Each test solution

(100 μ L, 5 replicates) produced a lower background signal (B) in the absence of AP and a higher signal (S) than the control when reacted with 4×10^{-16} mol of AP after a 3.5 min incubation at room temperature. In this experiment, 5 the maximum signal was obtained using 1.0 μ g/mL while the maximum S/B was obtained using 1.0 mg/mL.

Example 41. Western Blot Assay. A composition of the present invention were used to detect and quantify a 10 protein, human transferrin in a Western blot with an AP-labeled antibody on a polyvinylidene difluoride (PVDF) membrane. Dilutions of transferrin containing from 5000, 1000, 180, 30 and 5 pg, respectively, of protein were electrophoresed and transferred to PVDF membranes 15 (Millipore, Bedford, MA). The transferrin bands were blocked with and then reacted sequentially with goat anti-human transferrin and rabbit anti-goat-AP conjugate. The membranes were soaked briefly with reagents comprising 0.1 M tris buffer, pH 8.8 containing 0.88 mM $MgCl_2$, 0.66 mM 20 compound 5 and 64 μ M lucigenin. The membranes were placed between transparent plastic sheets and exposed to X-ray film. The bands for human transferrin were detected after 10 min with a 2.5 min exposure. The light produced using this composition led to intense emission which could be 25 imaged for several hours.

Similar results were obtained using nitrocellulose blotting membrane instead of PVDF membrane.

Example 42. Southern Blot Assay. A representative example of the use of a reagent of the present invention in a Southern blot assay is demonstrated in the following example.

- 5 Avidin-AP conjugate was obtained from Cappel Products (Durham, NC). Bovine serum albumin (heat shocked) was purchased from Sigma Chemical Co. (St. Louis, MO). Mouse genomic DNA was from Clontech (Palo Alto, CA). Biotin-ylated lambda DNA/HindIII fragments, biotin-7-dATP and the
- 10 Nick Translation kit were from Life Technologies, restriction endonuclease EcoRI was from Boehringer-Mannheim (Indianapolis, IN), and nylon membrane from Micron Separations Inc (Westborough, MA). X-ray film was X-OMAT AR from Kodak (Rochester, NY).
- 15 Mouse genomic DNA (30 µg) was cleaved with EcoRI to completion. Two and four µg portions were purified by phenol/chloroform extraction and ethanol precipitation. Purified DNA was separated by electrophoresis on 0.75% agarose gel with 40 mmol/L Tris-acetate, 2 mmol/L EDTA, pH
- 20 8.0 as the elution buffer. Following electrophoresis, the gel was rinsed with water, equilibrated for 12 min with 0.25 mol/L HCl, rinsed with water again, incubated in 0.5 mol/L NaOH, 1.5 mol/L NaCl for 15 min then in a fresh change of the same solution for 30 min, rinsed with water,
- 25 and incubated in three changes of 1 mol/L tris, 1.5 mol/L NaCl, pH 7.5 for 15 min each.

Nylon membrane was soaked sequentially with water for two min and 10X SSC for 30 min. DNA was transferred to the

membrane by capillary blotting overnight in 10X SSC. The membrane was washed by gentle agitation in 10X SSC for 10 min at room temperature and air-dried on Whatman 3MM blotting paper for 30 min and baked at 80° C under vacuum for 2 h.

The membrane was soaked in 6X SSPE (20X SSPE is 3 mol/L NaCl, 0.2 mol/L NaH₂PO₄, 20 mmol/L EDTA, pH 7.4) followed by prehybridization for 3 h at 42° C in pre-hybridization solution: 6X SSPE, 50% freshly deionized formamide, filtered 5X Denhardt's solution (50X is 1% Ficoll 400, 1% PVP, 1% BSA (initial fraction by heat shock)), filtered 1% SDS and 200 µg/mL sheared, denatured herring sperm DNA. The hybridization probe, oncogene probe v-mos DNA (PanVera Corp., Madison, WI) was labeled with biotin-7-dATP by nick translation according to the manufacturers instructions. Genomic DNA was hybridized overnight at 42° C in 6X SSPE, 45% formamide, filtered 5X Denhardt's, 1% SDS, 200 µg/mL herring sperm DNA with 500 ng of denatured biotinylated probe. The biotinylated probe was denatured by boiling 4 min and cooling 10 min at 0° C. The membrane was washed twice with 0.5X SSC, 0.4% SDS at 25° C for 15 and 35 min respectively, three times with 0.5X SSC, 0.4% SDS at 55 ° C for 10 min, once with 2X SSC at 25° C for 5 min, twice with TBS (50 mmol/L Tris-HCl, pH 7.4, 0.15 mol/L NaCl) for 3 min. After the wash step, the membrane was blocked at 65° C for 1 h in filtered 3% BSA, 100 mmol/L Tris-HCl, pH 7.4, 0.15 mol/L NaCl and washed in T-TBS (0.05% Tween 20 in TBS) for 1 min.

The blocked membrane was incubated with a 1:2000 dilution of avidin-AP in T-TBS for 12 min followed by four fresh changes of T-TBS for 5, 10, 15 and 20 min followed by a final wash with TBS for 5 min. Excess buffer was
5 drained off and blots soaked in either the detection reagent described in Example 41 for 4 min. Excess reagent was drained off, and the blots placed between transparent sheets and exposed to X-ray film.

The single copy gene was detected at 14.5 kbp by
10 soaking the membrane with detection reagent and exposure to x-ray film for varying lengths of time. After an incubation period of 9 min, a band corresponding to the single copy gene in both fractions was visible with a 5 min exposure to film. Multiple exposures could be
15 performed for a day by using longer exposures.

Example 43. Chemiluminescent Immunoassay of hCG

A method for detection of hCG by chemiluminescent immunoassay was performed on the IMMULITE Automated
20 Analyzer using an IMMULITE kit supplied by Diagnostic Products Corp. (Los Angeles, CA) according to the protocol provided by the manufacturer. Reagent A of Example 32 was substituted for the detection reagent supplied in the kit. The software was modified to allow a shorter substrate
25 incubation. All data points are the average of triplicate tests except for blank readings which were the average of 5 tests of the sample diluent. Analytes were prepared by serial dilution of special calibrators with the standard

diluents supplied in the kit. Chemiluminescence measurements were made at 1.5 minutes after substrate introduction.

A minor modification of the injection mechanism was made to minimize the dead volume of chemiluminescent reagent in the tubing. The substrate heater was disconnected and an 18" x 1/16" OD clear teflon tubing wrapped in black electrical tape having flared ends was connected directly between the substrate pump and the gray substrate dispensing nozzle. A similar length of the same teflon tubing was attached to the inlet port of the substrate pump. This inlet tubing was placed in the bottle containing the substrate. No attempt was made to preheat or temperature control the substrate. Insensitivity to temperature is indicated by the tight CV's even in the absence of external substrate temperature control.

Independent tests on rapid reading luminometers indicate that the substrate can be read with as little as 10 seconds incubation time yielding no loss of sensitivity or accuracy. Assay results as shown in Figure 13 and Table 6 demonstrate the utility of the present compositions in a sandwich-type immunoassay.

Table 6. IMMULITE hCG Assay Tabulated Data

	<u>mIU/mL hCG</u>	<u>Intensity (CPS)</u>	<u>% CV</u>
25	6675	47332040	3.8
	3000	29174720	1.2
	1000	11031867	1.5

	300	2906560	2.5
	100	977760	2.6
	30	288883	0.7
	10	133573	6.3
5	3	71507	5.9
	1	54738	7.4
	Blank	47634	9.0

Example 44. Rapid Chemiluminescent Immunoassay of TSH

10 A method for detection of TSH by chemiluminescent immunoassay was performed on the IMMULITE Automated Analyzer using an IMMULITE RTH Rapid TSH Assay kit from Diagnostic Products Corp. according to the manufacturer's protocol with modifications as described in Example 43.

15 Reagent A of Example 32 was substituted for the detection reagent supplied in the kit.

 The reagent background was tested by removing the antibody conjugates from a Third Generation TSH test wedge, carefully washing the wedge and replacing the contents with
20 Type 1 water. The observed substrate background was approximately 5000 counts per second. This indicates that the non-specific binding background is several times the reagent background. Therefore, it is possible that reoptimization of the biochemistry in the wedge could
25 significantly extend the sensitivity of several of the assays and/or allow a reduction of the biochemical incubation time. Chemiluminescence measurements were made at 1.5 minutes after substrate introduction.

Table 7. IMMULITE RTH Rapid TSH Assay Data

	<u>μIU/mL TSH</u>	<u>Intensity (CPS)</u>	<u>% CV</u>
	75	17370000	0.9
	10	2263413	5.3
5	1	202120	2.6
	0.3	72357	2.6
	0.1	36887	1.5
	0.03	25607	2.6
	0.01	22647	5.0
10	Blank	20810	2.0

Assay results as shown in Figure 14 and Table 7 demonstrate the utility of the present compositions in providing a rapid assay.

15 Example 45. Chemiluminescent Immunoassay of TSH

A method for detection of TSH by chemiluminescent immunoassay was performed on the IMMULITE Automated Analyzer using an IMMULITE TSH Third Generation TSH Assay kit from Diagnostic Products Corp. according to the manufacturer's protocol with modifications as described in Example 43. Reagent A of Example 32 was substituted for the detection reagent supplied in the kit. Chemiluminescence measurements were made at 1.5 minutes after substrate introduction.

Table 8. IMMULITE Third Generation TSH Assay Data

	<u>μIU/mL TSH</u>	<u>Intensity (CPS)</u>	<u>% CV</u>
	75	25249293	1.6
	10	4265600	1.5
5	1	415363	0.8
	0.3	122580	4.3
	0.1	54070	0.3
	0.03	23810	3.7
	0.01	17240	0.4
10	0.003	13900	1.7
	Blank	12996	4.4

The assay results as shown in Figure 15 and Table 8 demonstrate the utility of the present compositions in providing a highly sensitive assay.

15

Example 46. Chemiluminescent Immunoassay of Estradiol. A

method for detection of estradiol by chemiluminescent immunoassay was performed on the IMMULITE Automated Analyzer using a kit supplied by Diagnostic Products Corp.

20 according to the manufacturer's protocol with modifications as described in Example 43. Reagent A of Example 32 was substituted for the detection reagent supplied in the kit. Chemiluminescence measurements were made at 1.5 minutes after substrate introduction.

Table 9. IMMULITE E2 Estradiol Assay Data

	<u>pg/mL Estradiol</u>	<u>Intensity (CPS)</u>	<u>% CV</u>
	2000	500300	4.1
	1000	783877	3.3
5	300	2521788	2.7
	100	5398650	2.9
	30	8808133	1.8
	10	10036147	2.1
	Blank	10536776	3.9

- 10 The assay results as shown in Figure 16 and Table 9 demonstrate the utility of the present compositions in competitive-type immunoassays.

Example 47. Chemiluminescent Detection of Acid Phosphatase

- 15 (AcP). Reagent A of Example 32 was used in an experiment to detect acid phosphatase. Reaction of 100 μ L of this composition with AcP (Sigma AcP LIN-TROL, reconstituted to 2.0 mL, 83 U/L) at ambient temperature in a test tube housed in a Turner TD-20e luminometer produced
- 20 chemiluminescence which reached maximum intensity within 2-5 sec and decayed to zero within 10-15 sec.

Acid phosphatase could be measured in the presence of alkaline phosphatase by means of this detection reagent. Since the chemiluminescent signal induced by AcP decayed

25 nearly completely within seconds, measuring the stable light intensity after about 15-20 sec allows discrimination of AP activity as shown in Figure 17. This method allowed the simultaneous quantitation of both AcP and AP activity.

in the same sample in one experiment. AcP could be easily measured in human whole blood.

Example 48. Linearity and Sensitivity of Detection of AP

5 using Compound 13. In the manner of Example 32, a reagents was prepared according to the composition below. To 100 μL portions of each of the reagent was added 10 μL dilutions of AP containing between 8×10^{-16} mol and 8×10^{-22} mol of enzyme or 10 μL of water for the reagent blank. Light
10 intensity was measured at 75 sec. Figure 19 depicts the results.

Reagent

Compound 13, 0.33 mM

0.1 M tris buffer, pH 8.8

15 MgCl_2 , 5 μM

lucigenin, 3.2 μM

SDS, 0.5 mg/mL


Na_2SO_3 , 5 $\mu\text{g/mL}$

TWEEN 20, 0.15 mg/mL (w/v)

20


Example 49. The following compounds have also been synthesized and found to provide chemiluminescence when reacted with AP:

25



wherein V is t-butyl, CH₃, OCH₃, F, Cl, Br, I, COCH₃, CN and NO₂ as well as compounds of the formula:

5




wherein U is p-I, p-CH₃, m-OCH₃, o-Cl, m-Cl, o-Br, m-Br, p-Br and p-NO₂ as well as compounds of this formula having a 3,4-dichloro-, 2,5-dichloro- and 2,6-dichlorophenyl group.

The foregoing description and examples are illustrative only and not to be considered as restrictive. It is recognized that modifications of the specific compounds and methods not specifically disclosed can be made without departing from the spirit and scope of the present invention. The scope of the invention is limited only by the appended claims.

WHAT IS CLAIMED IS:

1. A compound of formula I:



Lumigen's Broadcast
Claims in WO 97/26245

I

5

wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality.


10

2. The compound of Claim 1 wherein R₆ contains from 1 to 50 atoms selected from the group consisting of C, N, O, S, P and halogen atoms.


3. The compound of Claim 2 wherein R₆ is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.

4. The compound of Claim 1 which is selected from the group consisting of compounds of formula II and compounds of formula III:

5



II




III

wherein R_1 is an organic group containing from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring.

5. The compound of Claim 4 wherein R_1 is selected from the group consisting of C_1 to C_4 alkyl groups and a benzyl group.


6. The compound of Claim 4 having the formula:



wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced and wherein Ar is an aryl ring group.

7. The compound of Claim 6 wherein at least one of R_7 to R_{14} is an alkoxy group and the remaining of R_7 to R_{14} are hydrogen.


8. The compound of Claim 6 having the formula:



wherein Ph is a phenyl group and each M is an alkali metal ion and n is 2.

9. The compound of Claim 8 wherein M is Na and n is 2.

10. The compound of Claim 6 having the formula:




wherein Ph is a phenyl group and each M is an alkali metal ion and n is 2.

11. The compound of Claim 10 wherein M is Na and n is 2.


12. A compound having the formula:

5




13. A compound having the formula:

5




14. A compound having the formula:

5




15. A compound having the formula:

5




16. A compound having the formula:

5




17. A compound having the formula:

5




18. A compound having the formula:

5



19. A reagent composition which produces chemiluminescence in the presence of a phosphatase enzyme which comprises in an aqueous solution:

a) a compound of formula I:




5

which reacts with the phosphatase enzyme wherein Het is a
10 heterocyclic ring system comprising at least one five or
six-membered ring which comprises at least one heteroatom
selected from the group consisting of N, O and S atoms,
wherein Z is selected from the group consisting of O and S
atoms, wherein R₆ is an organic group which allows
15 chemiluminescence to be produced, wherein each M is
independently selected from H and a cationic center and
wherein n is a number which satisfies electroneutrality;
and


b) at least one surfactant enhancer in an amount
20 effective to enhance the chemiluminescence.

20. The composition of Claim 19 in which the compound of formula I is selected from the group consisting of compounds of formula II and compounds of formula III:

5



II




III

wherein R_1 is an organic group selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring.

21. The composition of Claim 20 in which the compound of formula I has the formula:

5




wherein R_1 is selected from the group consisting of alkyl, heteroalkyl and aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced and wherein Ar is an aryl ring group.

22. The composition of Claim 21 wherein at least one of R_7 to R_{14} is an alkoxy group and the remaining of R_7 to R_{14} are hydrogen.

23. The composition of Claim 21 wherein the compound of formula I has the formula:


5



wherein Ph is a phenyl group.

24. The composition of Claim 21 wherein the compound has the formula:


5



wherein Ph is a phenyl group.

25. The composition of Claim 21 wherein the compound has the formula:

5



26. The composition of Claim 19 wherein the surfactant enhancer is selected from polymeric quaternary ammonium and phosphonium salts, monomeric quaternary ammonium and phosphonium salts and dicationic quaternary ammonium and
5 phosphonium salts.


27. The composition of Claim 19 wherein the surfactant enhancer is a copolymer of a vinylbenzyltributylphosphonium salt and a vinylbenzyltrioctylphosphonium salt.

28. The composition of Claim 19 wherein copolymer comprises vinylbenzyltributylphosphonium salt and vinylbenzyltrioctylphosphonium salt groups in a ratio of about 3 to 1.

29. The composition of Claim 19 further comprising a magnesium salt in an amount effective to promote the reaction.

30. A method for producing chemiluminescence which comprises reacting a phosphatase enzyme with at least one compound of formula I:

5




wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality.

10


15

31. The method of Claim 30 wherein the compound of formula I is selected from the group consisting of compounds of formula II and compounds of formula III:

5



II




III

wherein R_1 is an organic group containing from 1 to 50 atoms selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring.

32. The method of Claim 30 wherein the compound of formula I has the formula:

5




wherein R_1 is selected from the group consisting of alkyl, heteroalkyl and aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent which can contain from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced and wherein Ar is an aryl ring group.

33. The method of Claim 32 wherein at least one of R_7 to R_{14} is an alkoxy group and the remaining of R_7 to R_{14} are hydrogen.

34. The method of Claim 32 wherein the compound of formula I has the formula:


5



wherein Ph is a phenyl group.

35. The method of Claim 32 wherein the compound of formula I has the formula:


5



wherein Ph is a phenyl group.

36. The method of Claim 32 wherein the compound of formula I has the formula:

5



37. The method of Claim 30 wherein the phosphatase enzyme is selected from the group consisting of bacterial alkaline phosphatase, mammalian alkaline phosphatase, plant acid phosphatase, mammalian acid phosphatase and
5 alkaline phosphatase conjugates.

38. The method of Claim 37 wherein the alkaline phosphatase conjugate comprises alkaline phosphatase linked to a biological molecule selected from the group consisting of haptens, antibodies, proteins, nucleic acids
5 and oligonucleotides.

39. The method of Claim 32 wherein the phosphatase enzyme is selected from the group consisting of bacterial alkaline phosphatase, mammalian alkaline phosphatase, plant acid phosphatase, mammalian acid phosphatase and
5 alkaline phosphatase conjugates.


40. The method of Claim 39 wherein the alkaline phosphatase conjugate comprises alkaline phosphatase linked to a biological molecule selected from the group consisting of haptens, antibodies, proteins, nucleic acids
5 and oligonucleotides.

41. The method of any of Claims 34, 35 or 36 wherein the phosphatase enzyme is selected from the group consisting of bacterial alkaline phosphatase, mammalian alkaline phosphatase, plant acid phosphatase, mammalian acid
5 phosphatase and alkaline phosphatase conjugates.

42. The method of Claim 41 wherein the alkaline phosphatase conjugate comprises alkaline phosphatase
linked to a biological molecule selected from the group consisting of haptens, antibodies, proteins, nucleic acids
5 and oligonucleotides.

43. A method for detecting an analyte in a sample by a chemiluminescent assay procedure which comprises:

- (a) reacting a phosphatase enzyme with at least one compound of formula I to produce chemiluminescence for
5 detecting the analyte,




- 10 wherein in the compound of formula I, Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S
15 atoms, wherein R_6 is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality;


- (b) detecting the chemiluminescence; and
20 (c) relating the amount of the chemiluminescence to the amount of the analyte.

44. The method of Claim 43 wherein the compound of formula I is selected from the group consisting of compounds of formula II and compounds of formula III

5



II




III

wherein R_1 is an organic group containing from 1 to 50.
 10 atoms selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced
 15 and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring.

45. The method of Claim 43 wherein the compound of formula I has the formula:

5




wherein R_1 is selected from the group consisting of alkyl, heteroalkyl and aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced and wherein Ar is an aryl ring group.

46. The method of Claim 45 wherein at least one of R_7 to R_{14} is an alkoxy group and the remaining of R_7 to R_{14} are hydrogen.

47. The method of Claim 43 wherein the compound of formula I has the formula:


5



wherein Ph is a phenyl group.

48. The method of Claim 43 wherein the compound of formula I has the formula:


5



wherein Ph is a phenyl group.

49. The method of Claim 43 wherein the compound of formula I has the formula:

5



50. The method of Claim 43 wherein the analyte to be detected is the phosphatase enzyme.

51. The method of Claim 43 wherein the analyte to be detected is an inhibitor of the phosphatase enzyme.

52. The method of Claim 43 further comprising reacting the analyte in the sample with an analyte binding compound which specifically binds with the analyte wherein the
5 analyte-binding compound is labeled with alkaline phosphatase.

53. The method of Claim 52 wherein the analyte-binding compound is selected from the group consisting of antibodies, antigens, haptens and nucleic acids.

54. The method of Claim 45 further comprising reacting the analyte in the sample with an analyte binding compound which specifically binds with the analyte and at least one phosphatase-labeled specific binding substance for the
5 analyte binding compound.

55. The method of any of Claims 47, 48 or 49 further comprising reacting the analyte in the sample with an analyte binding compound which specifically binds with the analyte and at least one phosphatase-labeled specific
5 binding substance for the analyte binding compound.

56. The method of Claim 43 further comprising reacting the analyte in the sample with:

(a) a labeled analyte binding compound comprising an
5 analyte binding compound which specifically binds with the analyte and at least one second specific binding substance; and

(b) a phosphatase-labeled binding partner for the second specific binding substance.

57. The method of Claim 43 wherein the detection is performed on a membrane.

58. The method of Claim 57 wherein the membrane is selected from the group consisting of a nitrocellulose membrane, a polyvinylidene difluoride membrane and a nylon membrane.

59. The method of Claim 43 further comprising providing the compound of formula I in a reagent composition comprising, in an aqueous solution, the compound of formula I and a surfactant enhancer in an amount effective to enhance the chemiluminescence.

60. The method of Claim 45 further comprising providing the compound of formula I in a reagent composition comprising, in an aqueous solution, the compound of formula I and a surfactant enhancer in an amount effective to enhance the chemiluminescence.

61. The method of any of Claim 47, 48 or 49 further comprising providing the compound of formula I in a reagent composition comprising, in an aqueous solution, the compound of formula I and a surfactant enhancer in an amount effective to enhance the chemiluminescence.

62. The method of Claim 59 wherein the surfactant enhancer is a copolymer of a vinylbenzyltributylphosphonium salt and a vinylbenzyltrioctylphosphonium salt.

63. The method of Claim 43 further comprising:

(a) reacting the compound of formula I with the phosphatase enzyme in a buffer at a first pH for a first period of time;

5 (b) adding a strongly basic trigger solution to the buffer solution to raise the pH of the buffer to a second pH for inducing the chemiluminescence; and

(c) measuring the chemiluminescence.

64. The method of Claim 63 wherein the first pH is in the range 5.0 - 9.5, wherein the pH of the trigger solution is greater than about 11 and the first period of time is from about 1 seconds to about 10 minutes.


65. The method of Claim 63 wherein the basic trigger solution contains a surfactant enhancer.

66. The method of Claim 65 wherein the enhancer is a copolymer of a vinylbenzyltributylphosphonium salt and a vinylbenzyltrioctylphosphonium salt.

67. A reagent composition which produces chemiluminescence in the presence of a phosphatase enzyme which comprises in an aqueous solution:

a) a compound of formula I:

5



10 which reacts with the phosphatase enzyme wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S


15 atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality; and

20 b) a cationic aromatic compound in an amount effective to increase the chemiluminescence compared to that generated in the absence of the cationic aromatic compound.

68. The composition of Claim 67 wherein R_6 contains from 1 to 50 atoms selected from the group consisting of C, N, O, S, P and halogen atoms.

69. The composition of Claim 68 wherein R_6 is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.


70. The composition of Claim 67 in which the compound of formula I is selected from the group consisting of compounds of formulas:



wherein R_1 is an organic group containing from 1 to 50
 10 atoms selected from C, N, O, S, P and halogen atoms and is
 selected from the group consisting of alkyl, substituted
 alkyl, aryl, substituted aryl, aralkyl and substituted
 aralkyl groups, each of R_2 to R_5 is a substituent
 containing from 1 to 50 atoms selected from C, H, N, O, S,
 15 P and halogen atoms which permits the chemiluminescence to
 be produced and wherein pairs of adjacent groups can be
 joined together to form a carbocyclic or heterocyclic ring
 system comprising at least one 5 or 6-membered ring.


71. The composition of Claim 67 in which the compound of formula I has the formula:

5




wherein R_1 is selected from the group consisting of substituted and unsubstituted C_1 to C_4 alkyl groups and substituted and unsubstituted aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced.

72. The composition of Claim 71 wherein the compound of formula I has the formula:



73. The composition of Claim 72 wherein the compound of formula I has the formula:



74. The composition of any one of Claims 67, 68, 69, 70, 71, 72, or 73 wherein the cationic aromatic compound is selected from the group consisting of cyanine dyes, carbocyanine dyes, azo dyes, acridinium derivatives, methylene blue, Nile Blue, IR-1040, lucigenin and paraquat dichloride.

75. The composition of Claim 67 further comprising an anionic surfactant in an amount effective to increase the speed with which maximum chemiluminescence intensity is reached and a non-ionic surfactant in an amount effective to increase the amount of chemiluminescence.

76. The composition of Claim 75 wherein the anionic surfactant is selected from alkylsulfates containing at least 10 carbon atoms and alkylsulfonates containing at least 10 carbon atoms.

77. The composition of Claim 75 wherein the non-ionic surfactant is selected from polyoxyethylenated alkylphenols, polyoxyethylenated alcohols, polyoxyethylenated ethers and polyoxyethylenated sorbitol esters.


78. The composition of Claim 75 additionally comprising a sulfite salt in an amount effective to reduce chemiluminescence produced by the composition in the absence of a phosphatase enzyme.

79. The composition of Claim 76 wherein the anionic surfactant is sodium dodecyl sulfate.

80. The composition of Claim 78 wherein the sulfite salt is sodium sulfite.

81. A method for producing chemiluminescence which comprises reacting a phosphatase enzyme with

a) at least one compound of formula I:




wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality; and

b) a cationic aromatic compound in an amount effective to increase the chemiluminescence compared to that generated in the absence of the cationic aromatic compound.

82. The method of Claim 81 wherein R_6 contains from 1 to 50 atoms selected from the group consisting of C, N, O, S, P and halogen atoms.

83. The method of Claim 82 wherein R_6 is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.


84. The method of Claim 81 wherein the compound of formula I is selected from the group consisting of compounds of formulas:



wherein R_1 is an organic group containing from 1 to 50
 10 atoms selected from C, N, O, S, P and halogen atoms and is
 selected from the group consisting of alkyl, substituted
 alkyl, aryl, substituted aryl, aralkyl and substituted
 aralkyl groups, each of R_2 to R_5 is a substituent
 containing from 1 to 50 atoms selected from C, H, N, O, S,
 15 P and halogen atoms which permits the chemiluminescence to
 be produced and wherein pairs of adjacent groups can be
 joined together to form a carbocyclic or heterocyclic ring
 system comprising at least one 5 or 6-membered ring.


85. The method of Claim 84 wherein the compound of formula I has the formula:

5




wherein R_1 is selected from the group consisting of substituted and unsubstituted C_1 to C_4 alkyl groups and substituted and unsubstituted aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent which can contain from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced.

86. The method of Claim 85 wherein the compound of formula I has the formula:



87. The method of Claim 86 wherein the compound of formula I has the formula:



88. The method of any of Claims 81, 82, 83, 84, 85, 86 or 87 wherein the phosphatase enzyme is selected from the group consisting of bacterial alkaline phosphatase, mammalian alkaline phosphatase, plant acid phosphatase, mammalian acid phosphatase and alkaline phosphatase conjugates.


5

89. The method of Claim 88 wherein the alkaline
phosphatase conjugate comprises alkaline phosphatase
linked to a biological molecule selected from the group
consisting of haptens, antibodies, proteins, nucleic acids
5 and oligonucleotides.

90. The method of any of Claims 81, 82, 83, 84, 85, 86 or
87 further comprising providing the compound of formula I
and the cationic aromatic compound in a reagent
5 composition which composition further comprises, in an
aqueous solution, an anionic surfactant in an amount
effective to increase the speed with which maximum
chemiluminescence intensity is reached and a non-ionic
surfactant in an amount effective to increase the
10 chemiluminescence.

91. A method for detecting an analyte in a sample by a chemiluminescent assay procedure which comprises:

- (a) reacting a phosphatase enzyme with at least one compound of formula I in the presence of a cationic aromatic compound to produce chemiluminescence for detecting the analyte,



10

wherein in the compound of formula I, Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality;

20


(b) detecting the chemiluminescence; and

(c) relating the amount of the chemiluminescence to the amount of the analyte.

92. The method of Claim 91 wherein R_6 contains from 1 to 50 atoms selected from the group consisting of C, N, O, S, P and halogen atoms.

93. The method of Claim 92 wherein R_6 is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.

94. The method of Claim 91 wherein the compound of formula I is selected from the group consisting of compounds of formulas:




wherein R_1 is an organic group containing from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, aralkyl and substituted aralkyl groups, each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring.

10

15


95. The method of Claim 91 wherein the compound of formula I has the formula:

5




wherein R_1 is selected from the group consisting of substituted and unsubstituted C_1 to C_4 alkyl groups and substituted and unsubstituted aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced.

96. The method of Claim 95 wherein the compound of formula I has the formula:



5

97. The method of Claim 96 wherein the compound of formula I has the formula:



98. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 wherein the analyte to be detected is the phosphatase enzyme.

99. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 wherein the analyte to be detected is an inhibitor of the phosphatase enzyme.


100. The method of any one of Claims 91,
96 or 97 further comprising reacting the a:
sample with a labeled analyte binding compo
specifically binds with the analyte wherein
5 analyte-binding compound is labeled with a
phosphatase.

101. The method of Claim 100 wherein the a
compound is selected from the group consist
antibodies, antigens, haptens and nucleic a

102. The method of any one of Claims 91, 9
96 or 97 further comprising reacting the ar
sample with an analyte binding compound whi
binds with the analyte and at least one phc
5 labeled specific binding substance for the
compound.

95. The method of Claim 91 wherein the compound of formula I has the formula:


5




wherein R_1 is selected from the group consisting of substituted and unsubstituted C_1 to C_4 alkyl groups and substituted and unsubstituted aralkyl groups, wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced.

96. The method of Claim 95 wherein the compound of formula I has the formula:

5



97. The method of Claim 96 wherein the compound of formula I has the formula:



98. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 wherein the analyte to be detected is the phosphatase enzyme.

99. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 wherein the analyte to be detected is an inhibitor of the phosphatase enzyme.

100. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 further comprising reacting the analyte in the sample with a labeled analyte binding compound which specifically binds with the analyte wherein the
5 analyte-binding compound is labeled with alkaline phosphatase.

101. The method of Claim 100 wherein the analyte-binding compound is selected from the group consisting of antibodies, antigens, haptens and nucleic acids.

102. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 further comprising reacting the analyte in the sample with an analyte binding compound which specifically binds with the analyte and at least one phosphatase-
5 labeled specific binding substance for the analyte binding compound.

103. The method of any one of Claims 91, 92, 93, 94, 95, 96 or 97 further comprising reacting the analyte in the sample with:

(a) a labeled analyte binding compound comprising an
5 analyte binding compound which specifically binds with the analyte and at least one second specific binding substance; and

(b) a phosphatase-labeled binding partner for the second specific binding substance.


104. The method of Claim 91 wherein the detection is performed on a membrane.

105. The method of Claim 104 wherein the membrane is selected from the group consisting of a nitrocellulose membrane, a polyvinylidene difluoride membrane and a nylon membrane.

106. The method of any one of Claims 91, 92, 93, 94, 95,
96 or 97 further comprising providing the compound of
formula I and the cationic aromatic compound in a reagent
composition which composition further comprises, in an
5 aqueous solution, an anionic surfactant in an amount
effective to increase the speed with which maximum
chemiluminescence intensity is reached and a non-ionic
surfactant in an amount effective to increase the
chemiluminescence.

107. A method for detecting acid phosphatase and alkaline phosphatase in a sample suspected of containing both acid phosphatase and alkaline phosphatase by a chemiluminescent assay procedure which comprises:

- 5 (a) reacting the sample with a reagent composition comprising at least one compound of formula I,



10

- wherein in the compound of formula I, Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms,
- 15 wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality,
- 20 a cationic aromatic compound, an anionic surfactant in an amount effective to increase the speed with which maximum chemiluminescence intensity is reached and a non-ionic surfactant in an amount effective to increase the chemiluminescence;

(b) detecting the amount or intensity of chemiluminescence during an initial period;


(c) waiting a second period of time until the chemiluminescence has achieved a constant level;

5 (d) detecting the amount or intensity of chemiluminescence during a third period;

(e) relating the chemiluminescence in the initial time period to the amount of acid phosphatase; and

(f) relating the chemiluminescence in the third time
10 period to the amount of alkaline phosphatase.


108. The method of Claim 107 wherein the compound of formula I is selected from the group consisting of compounds of formulas:



wherein R_1 is an organic group containing from 1 to 50
 10 atoms selected from C, N, O, S, P and halogen atoms and is
 selected from the group consisting of alkyl, substituted
 alkyl, aryl, substituted aryl, aralkyl and substituted
 aralkyl groups, each of R_2 to R_5 is a substituent
 containing from 1 to 50 atoms selected from C, H, N, O, S,
 15 P and halogen atoms which permits the chemiluminescence to
 be produced and wherein pairs of adjacent groups can be
 joined together to form a carbocyclic or heterocyclic ring
 system comprising at least one 5 or 6-membered ring.

109. The method of Claim 107 wherein the compound of formula I has the formula:

5




wherein R₁ is selected from the group consisting of substituted and unsubstituted C₁ to C₄ alkyl groups and substituted and unsubstituted aralkyl groups, wherein each of R₇ to R₁₄ is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced.


10

110. The method of Claim 109 wherein the compound of formula I has the formula:

5



111. The method of Claim 107 wherein the compound of formula I has the formula:



112. A process for the preparation of a compound of formula I:



wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one heteroatom selected from the group consisting of N, O and S atoms, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced, wherein each M is independently selected from H and a cationic center and wherein n is a number which satisfies electroneutrality comprising the steps of:

10


15

(a) reacting a heterocyclic ester or thioester compound VIII having the formula:



wherein Het, Z and R₆ are as defined for compound I with a base to form an enolate of VIII;

(b) reacting the enolate with a phosphorylating agent to form a protected enol phosphate IX having the formula:



wherein Het, Z and R₆ are as defined above and Y is a


5 protecting group; and

(c) deprotecting the enol phosphate to form the enol phosphate salt compound I by reacting IX with at least one deprotecting agent in the presence of a cationic species M if the cationic species is not a part of the deprotecting

10 agent.

113. The process of Claim 112 wherein the step of reacting the enolate of compound VIII with a phosphorylating agent to form a protected enol phosphate IX comprises the steps of :

- 5 (a) reacting the enolate of compound VIII with a phosphorus oxyhalide compound POW_3 , where W is a halogen atom selected from F, Cl, Br and I to form an enol dihalophosphate X having the formula



wherein Het, Z and R_6 are as defined in compound VIII; and

- (b) reacting compound X with at least two equivalents
 15 of a hydroxylic compound Y-OH to form the protected enol phosphate IX.

114. The process of Claim 112 wherein the step of reacting the enolate of compound VIII with a phosphorylating agent to form a protected enol phosphate IX comprises reacting the enolate of compound VIII with a phosphorylating agent
 5 containing the protecting groups Y and having the formula W-PO(OY)_2 and wherein W is a halogen atom selected from F, Cl, Br and I.

115. The process of Claim 112 wherein the groups Y are selected from the group consisting of lower alkyl groups, substituted lower alkyl groups, phenyl, substituted phenyl and benzyl groups.

116. The process of Claim 112 wherein the groups Y are connected to form the single group $-\text{CH}_2\text{CH}_2-$.


117. The process of Claim 112 wherein the deprotecting agent is selected from the group consisting of organic and inorganic bases such as sodium hydroxide, potassium hydroxide, potassium carbonate, sodium methoxide, sodium
5 ethoxide, potassium t-butoxide, ammonium hydroxide, nucleophilic agents such as cyanide ion, fluoride ion.

118. The process of Claim 112 wherein Y is a $\text{CH}_2\text{CH}_2\text{CN}$ group and the deprotecting agent is selected from sodium hydroxide and sodium carbonate.


119. The process of Claim 113 wherein Y is a $\text{CH}_2\text{CH}_2\text{CN}$ group and the deprotecting agent is selected from sodium hydroxide and sodium carbonate.

120. The process of Claim 112 wherein the compound of formula I is selected from the group consisting of compounds of formula II and compounds of formula III:

5



II




III

wherein R_1 is an organic group containing from 1 to 50 •
 10 atoms selected from C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, wherein each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms
 15 which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring and wherein R_6 contains from 1 to 50 atoms selected from the group consisting of
 20 C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.

121. The process of Claim 120 wherein the compound of formula I has the formula:


5



wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced and wherein Ar is an aryl ring group.

10


122. A compound of formula IX:



5

wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one N or S atom, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an
10 organic group which allows chemiluminescence to be produced and wherein Y is a protecting group which is removable to form a phosphate salt compound.

123. The compound of Claim 122 which is selected from the group consisting of compounds of formulas:




wherein R_1 is an organic group containing from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, wherein each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring and wherein R_6 contains from 1 to 50 atoms selected from the group consisting of C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.

10

15

20

124. The compound of Claim 123 having the formula:




5

wherein each of R_7 to R_{14} is independently a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms and which permits the chemiluminescence to be produced and wherein Ar is an aryl ring group.

10

125. The compound of any of Claims 122, 123 or 124 wherein Y is a $\text{CH}_2\text{CH}_2\text{CN}$ group.

126. A compound of formula X:




5

wherein Het is a heterocyclic ring system comprising at least one five or six-membered ring which comprises at least one N or S atom, wherein Z is selected from the group consisting of O and S atoms, wherein R₆ is an organic group which allows chemiluminescence to be produced and wherein W is a halogen atom selected from F, Cl, Br and I atoms.

10

127. The compound of Claim 126 which is selected from the group consisting of compounds of formulas:




wherein R_1 is an organic group containing from 1 to 50 atoms selected from C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups, wherein each of R_2 to R_5 is a substituent containing from 1 to 50 atoms selected from C, H, N, O, S, P and halogen atoms which permits the chemiluminescence to be produced and wherein pairs of adjacent groups can be joined together to form a carbocyclic or heterocyclic ring system comprising at least one 5 or 6-membered ring and wherein R_6 contains from 1 to 50 atoms selected from the group consisting of C, N, O, S, P and halogen atoms and is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl and aralkyl groups.

10

15

20

128. The compound of Claim 127 having the formula:




5
 wherein each of R₇ to R₁₄ is independently a substituent
 containing from 1 to 50 atoms selected from C, H, N, O, S,
 P and halogen atoms and which permits the
 chemiluminescence to be produced and wherein Ar is an aryl
 10 ring group.

129. The compound of any of Claims 126, 127 or 128 wherein
 W is a Cl atom.


1/19

FIG. 1




2/19

FIG. 2




3/19

FIG. 3




4/19

FIG. 4



5/19

FIG. 5




6/19

FIG. 6




7/19

FIG. 7




8/19

FIG. 8




9/19

FIG. 9




10/19

FIG. 10




11/19

FIG. 11




12/19

FIG. 12




13/19

FIG. 13




14/19

FIG. 14




15/19

FIG. 15




16/19

FIG. 16




17/19

FIG. 17




18/19

FIG. 18



19/19

FIG. 19



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/00015

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07D 219/00

US CL : 546/23

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/23

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE, REGISTRY

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3,130,203 A (K.H. BUCHEL ET AL.) 21 April 1964, see compound 11.	1-3, 122
A	US 5,306,621 A (L.J. KRICKA) 26 April 1994, see entire document.	1-129
A	KITAMURA et al. A New Highly Sensitive Chemiluminescent Assay of Alkaline Phosphatase Using Lucigenin and its Application to Enzyme Immunoassay. J. Biolumin. Chemilumin. 1995, Vol. 10, pages 1-7, see entire document.	1-129
A	SASAMOTO et al. Chemiluminescent Assay of Alkaline Phosphatase Using Phenacyl Phosphate. Anal. Chim. Acta. 1995, Vol. 306, pages 161-166, see entire document.	1-129



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understate the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E earlier document published on or after the international filing date	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G	document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means		
*P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

12 MARCH 1997

Date of mailing of the international search report

27 MAR 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

GARTH M. DAHLEN

Facsimile No. (703) 305-3230

Telephone No. (703) 308-1235